UNIVERSIDAD DE CONCEPCIÓN



Centro de Investigación en Ingeniería Matemática (CI^2MA)



Numerical solution of a spatio-temporal predator-prey model with infected prey

> RAIMUND BÜRGER, GERARDO CHOWELL, Elvis Gavilán, Pep Mulet, Luis M. Villada

> > PREPRINT 2018-17

SERIE DE PRE-PUBLICACIONES

NUMERICAL SOLUTION OF A SPATIO-TEMPORAL PREDATOR-PREY MODEL WITH INFECTED PREY

RAIMUND BÜRGER

CI²MA and Departamento de Ingeniería Matemática, Universidad de Concepción, Casilla 160-C, Concepción, Chile

Gerardo Chowell

School of Public Health, Georgia State University, Atlanta, Georgia, USA Simon A. Levin Mathematical and Computational Modeling Sciences Center, School of Human Evolution and Social Change, Arizona State University, Tempe, AZ 85287, USA, and Division of International Epidemiology and Population Studies, Fogarty International Center, National Institutes of Health, Bethesda, MD 20892, USA

Elvis Gavilán

CI²MA and Departamento de Ingeniería Matemática, Universidad de Concepción, Casilla 160-C, Concepción, Chile

Pep Mulet

Departament de Matemàtiques, Universitat de València, Av. Dr. Moliner 50, E-46100 Burjassot, Spain

LUIS M. VILLADA

GIMNAP-Departamento de Matemáticas, Universidad del Bío-Bío, Casilla 5-C, Concepción, Chile CI²MA, Universidad de Concepción, Casilla 160-C, Concepción, Chile

(Communicated by the associate editor name)

ABSTRACT. A spatio-temporal eco-epidemiological model is formulated by combining an available non-spatial model for predator-prey dynamics with infected prey [D. Greenhalgh and M. Haque, Math. Meth. Appl. Sci., 30 (2007), 911-929] with a spatio-temporal susceptible-infective (SI)-type epidemic model of pattern formation due to diffusion [G.-Q. Sun, Nonlinear Dynamics, 69 (2012), 1097-1104]. It is assumed that predators exclusively eat infected prey, in agreement with the hypothesis that the infection weakens the prey and increases its susceptibility to predation. Furthermore, the movement of predators is described by a non-local convolution of the density of infected prey as proposed in [R.M. Colombo and E. Rossi, Commun. Math. Sci., 13 (2015), 369-400]. The resulting convection-diffusion-reaction system of three partial differential equations for the densities of susceptible and infected prey and predators is solved by an efficient method that combines weighted essentially non-oscillatory (WENO) reconstructions and an implicit-explicit Runge-Kutta (IMEX-RK) method for time stepping. Numerical examples illustrate the formation of spatial patterns involving all three species. Future directions of research are suggested.

Date: April 10, 2018.

²⁰¹⁰ Mathematics Subject Classification. Primary: 92D30, Secondary: 97M60, 65L06, 65M20. Key words and phrases. Convection-diffusion-reaction system, implicit-explicit Runge-Kutta scheme, nonlocal velocity, predator-prey model.

1. Introduction.

 $\mathbf{2}$

1.1. Scope. Mathematical models are powerful tools to characterize the spatialtemporal changes in complex dynamical systems. More specifically, mathematical models are frequently used to gain novel insight into the dynamics and control of natural and social phenomena [38]. For example, systems of mathematical models are often simple and crude approximations that aim to capture key mechanisms underlying the dynamics of infectious disease transmission in human, animal, and plant populations. Such first approximation models can be subsequently refined to study the effects of additional factors and processes including spatial heterogeneity in host density and their characteristics (e.g., fitness, susceptibility, infectivity) as well as specific host behaviors (e.g., movement patterns) in response to local host density or disease prevalence.

It is the purpose of this contribution to advance a spatio-temporal predatorprey model taking into account infected prey with a particular focus on its efficient numerical solution. The resulting eco-epidemiological model combines several submodels that have been proposed in the recent literature. The population of prey is described by a spatio-temporal SIR-type epidemiological model similar to the one studied by Sun [39] while the predator-prey interaction follows the treatment by Greenhalgh and Haque [20] (formulated in a non-spatial setting in that paper). The movement of predators is based on the nonlocal velocity model advanced for the predator movement in a spatio-temporal predator-prey model by Colombo and Rossi [13]. That velocity model was also used for the male compartments in a recently published model of hantavirus infection [12]. In particular the implicitexplicit numerical technique for efficiently solving the governing partial differential equations (PDEs) is adapted to the new model presented herein.

In most general terms, the governing model considers a prey population with a susceptible and infective compartment, denoted by S and I, and a variable Ydenoting the population of the predator. In the spatio-temporal context, these variables are understood as local densities that are functions of position $\boldsymbol{x} \in \Omega$ and time $t \in \mathcal{T} := [0, T]$ on a bounded domain $\Omega \subset \mathbb{R}^2$. The final model for

$$u := (u_1, u_2, u_3)^{\mathrm{T}} := (S, I, Y)^{\mathrm{T}}, \quad u = u(x, t)$$

is given by a convection-diffusion reaction system of the type

$$\frac{\partial \boldsymbol{u}}{\partial t} + \nabla \cdot \boldsymbol{F}^{c}(\boldsymbol{u}) = \boldsymbol{\mathcal{D}} \Delta \boldsymbol{u} + \boldsymbol{s}(\boldsymbol{u}), \qquad (1.1)$$

supplied with initial and boundary conditions, where the convective fluxes $F^{c}(u)$, the diffusion matrix \mathcal{D} , and the vector of reaction terms s(u) are specified in later parts of the paper.

It is proposed to describe the movement of the predators in a particular way that depends non-locally on the density I of infected prey, while the movement of infected and susceptible prey, that is of members of the compartments I and S, is standard diffusion. This property calls for numerical methods that allow the efficient computation of numerical fluxes based on the non-local evaluation of data but avoid the severe time step restriction incurred by explicit time discretizations of the diffusion term $\mathcal{D}\Delta u$. As in [12] the first goal can be achieved by a technique based on Fast Fourier Transforms (FFT) in combination with an implicit-explicit (IMEX) discretization to handle the second. The numerical method constructed in this way is applied to simulate several scenarios that allow comparing the effect of initial conditions and different parameter values. The numerical results exhibit a rich variety of behavior, including formation of patterns taking the shapes of spots, stripes, both from randomized perturbations of steady states and in a scenario that describes the invasion of a prey habitat by the predator. The structure of solutions obtained in the latter case are reminiscent of permanent wave fronts.

1.2. Related work. The spatial spread of infectious diseases is described mathematically in a number of monographs that include [15, 30, 31, 35, 42]. The temporal evolution of diseases is also treated in [10, 18]. Our assumption that all epidemiological compartments are distributed over the whole spatial domain is opposed to the alternative metapopulation approach that describes spatial structure through wellidentified sub-populations or "patches" (cf., e.g., [1-3, 11, 40, 41]). The description of spatial structure by explicitly specifying the mobilities between "patches" is typical for characterizing the behavior of humans who undertake directed travels, while a description through a convection-diffusion-reaction mechanism is more suitable for non-human infectious agents such as spores, insects, and bacteria that would disperse [15, 31, 32]. A decisive advantage of the spatially continuous approach, namely its amenability to mathematical analysis is emphasized in [43].

The introduction of [20] and Hadeler and Freedman [21] provide extensive references to real-world examples of three-species eco-epidemiological systems of sound prey, infected prey, and predators. Specifically, in [21] a predator-prey model is described in which the prey is infected by a parasite and the prey in turn infects the predator with that parasite. The models studied in [20] and herein are based on the general assumption that the infection weakens the prey and increases its susceptibility to predation.

A less common ingredient in mathematical epidemiology is the convective term $\nabla \cdot F^{c}(\boldsymbol{u})$. Related advection terms, for which the essential functional dependence for one compartment is $F^{c} = F^{c}(\boldsymbol{x}, \boldsymbol{u}) = \boldsymbol{b}(\boldsymbol{x})\boldsymbol{u}$ with a given velocity $\boldsymbol{b}(\boldsymbol{x})$, arise if the population under study is transported (as is the case with wind-borne infectious agents, plankton, etc.). In our model the main role of the convective term is similar to that of [31, Ch. 14] in that it imposes a preferred direction of movement of predators, where the global dependence of the direction is determined by convolution with population data within a certain horizon of the current position. This idea of non-local dependence of biological fluxes goes back to a non-local predator-prey model introduced and analyzed by Colombo and Rossi [13], and for which a numerical scheme was analyzed in [34]. The scheme of [34] is based on the Lax-Friedrichs scheme for the ease of demonstrating convergence properties; we here employ higher-order weighted essentially non-oscillatory (WENO) reconstructions [22, 29] to achieve high-order spatial accuracy.

Furthermore, IMEX Runge-Kutta (IMEX-RK) schemes play an important role for the efficient numerical solution of (1.1). Roughly speaking, an IMEX-RK method for the convection-diffusion-reaction equation (1.1) consists of a Runge-Kutta scheme with an implicit discretization of the diffusive term $D\Delta u$ combined with an explicit one for the convective and reactive terms, $F^{c}(u)$ and s(u), respectively. To introduce the main idea, we assume that

$$\frac{\mathrm{d}\boldsymbol{v}}{\mathrm{d}t} = \Phi^*(\boldsymbol{v}) + \Phi(\boldsymbol{v}) \tag{1.2}$$

represents a method-of-lines semi-discretization of (1.1), where $\Phi^*(v)$ and $\Phi(v)$ are spatial discretizations of the convective and reactive terms and of the diffusive term,

respectively. Assume that the spatial mesh width is h > 0 in both the x- and ydirection. Then the stability restriction on the time step Δt that explicit schemes impose when applied to (1.2) is very severe (Δt must be proportional to h^2), due to the presence of $\Phi(\mathbf{v})$. The implicit treatment of both $\Phi^*(\mathbf{v})$ and $\Phi(\mathbf{v})$ would remove any stability restriction on Δt . However, the upwind nonlinear discretization of the convective terms in $\Phi^*(\mathbf{v})$ that is needed for stability makes its implicit treatment extremely involved. This situation becomes even more complicated due to the use of WENO reconstructions [22, 29, 36, 44]. However, numerical integrators that deal implicitly with $\Phi(\mathbf{v})$ and explicitly with $\Phi^*(\mathbf{v})$ can be used with a time step restriction dictated by the convective-reactive term alone. These schemes have been profusely used in convection-diffusion problems and convection problems with stiff reaction terms [4,6–9,14,16,24,33,45].

1.3. **Outline of the paper.** The remainder of the paper is organized as follows. In Section 2 the governing mathematical model is introduced. To this end, we first formulate (in Section 2.1) a non-spatial predator-prev model, defined by three nonlinear ordinary differential equations (ODEs), in which an epidemic spreads in the prev. This model gives rise to five equilibrium points whose respective stability is discussed in Section 2.2. (These stability results will motivate the choice of parameters of the simulated spatio-temporal scenarios.) Then, in Section 2.3, we obtain the full spatio-temporal model by equipping the ODE model with diffusive terms for the prey species and a non-local convection term for predators. These ingredients are specified in detail in Section 4.3. Section 4 is devoted to the description of the numerical scheme proposed to solve the spatio-temporal model developed in Section 2.3. Specifically, the (standard) spatial discretization of local convection and diffusion terms is introduced in Section 3.1. The spatial discretization of the convolution defining the non-local predator velocities is described in Section 3.2. These velocities, in turn, are utilized in the definition of the final discretization of the convective flux by means of a fifth-order WENO discretization, as is detailed in Section 3.3. The time discretization of the complete model by an IMEX-RK scheme is outlined in Section 3.4. In Section 4, which is at the core of this paper, numerical solutions to the spatio-temporal model are presented. To motivate the choice of parameters, we focus on two equilibrium points of the three-equation ODE model of Section 2.1, as is briefly discussed in Section 4.1. In Section 4.2 we present a total number of 10 examples of numerical solutions of the spatio-temporal model that give rise to patterns. In Section 4.3 we present Example 11 that has been designed to demonstrate in a purely visual way the convergence of the numerical scheme to a definite solution upon refinement of the computational grid, while Example 12 illustrates the influence of the choice of the radius of the convolution kernel of the non-local prey velocity. Finally, some conclusions and possible directions of future work are collected in Section 5.

2. Mathematical model.

2.1. Dynamical system for a predator-prey model. Combining the SIR epidemic model of [39] with a second species corresponding to a predator (as in the model of [20]), and assuming that the predator only eats infected prey, we obtain the following model:

$$\frac{\mathrm{d}S}{\mathrm{d}t} = A - dS - \beta S I^2,\tag{2.1a}$$

4

$$\frac{\mathrm{d}I}{\mathrm{d}t} = \beta SI^2 - (d+\mu)I - \frac{cYI}{mY+I},\tag{2.1b}$$

$$\frac{\mathrm{d}Y}{\mathrm{d}t} = \delta Y \left(1 - \frac{hY}{I} \right),\tag{2.1c}$$

where t is time, S(t) is the population of susceptible prey, I(t) is the population of infected prey, A is the recruitment rate of the prey population, d is the natural death rate of the prey population, β is the force of infection or the rate of transmission, μ is the disease-related death from the infected, Y(t) is the population of predators, c is the search rate of the predators towards infected prey, δ is the per capita growth rate of the predators, h is a constant related to the density dependent mortality of the predator population, and m > 0 is a constant.

The incidence rate in (2.1a) and (2.1b) is chosen here as the nonlinear expression βSI^2 in agreement with Sun [39], who in turn appeals to the justification by Liu et al. [27,28]. The last term in the right-hand side of (2.1b) is a ratio-dependent predation term; see, e.g., [25] for further justification of such expressions. Furthermore, note that (2.1c) coincides with the predator equation [20, Eq. (1)], and that hY/I in that equation is the Leslie-Gower term [26] which measures the loss in the predator population due to the relative scarcity of the (infected) prey [19].

It is well known that the use of a ratio-dependent terms requires carefully handling situations of zero denominators. The right-hand sides of the ODE version (2.1) are potentially ill-defined if I(0) = 0. This issue is treated in [20] as follows: if I(0) = 0 and $Y(0) \ge 0$, then (2.1c) is interpreted as implying that Y(t) = 0 for t > 0 and if I(0) = Y(0) = 0, then (2.1b) is interpreted as implying that I(t) = 0 for all t. With respect to the PDE version (2.5), we have not encountered any difficulty of division by zero in our numerical experiments since in all cases we choose the initial datum $I(\mathbf{x}, 0) > 0$ for $\mathbf{x} \in \Omega$.

Theorem 2.1. The system (2.1) has the following equilibrium points:

- i) The equilibrium $E_1 = (A/d, 0, 0)$ corresponding to the extinction of the epidemic.
- ii) The following equilibria in absence of the predator:

$$E_2 = \left(\frac{A\beta + \sqrt{R}}{2d\beta}, \frac{2d(d+\mu)}{A\beta + \sqrt{R}}, 0\right), \quad E_3 = \left(\frac{A\beta - \sqrt{R}}{2d\beta}, \frac{2d(d+\mu)}{A\beta - \sqrt{R}}, 0\right),$$

where we define $R := A^2\beta^2 - 4d^3\beta - 8d^2\beta\mu - 4d\beta\mu^2$. These equilibria are feasible if R > 0, which occurs if $\beta > 4d(d + \mu)^2/A^2$.

iii) The following equilibria with presence of the predator:

$$E_4 = \left(\frac{A}{d + \beta I_4^2}, I_4, \frac{I_4}{h}\right), \quad E_5 = \left(\frac{A}{d + \beta I_5^2}, I_5, \frac{I_5}{h}\right), \tag{2.2}$$

where $I_4 < I_5$ are the solutions of the quadratic equation

$$\alpha\beta I^2 - \beta AI + d\alpha = 0$$
, where $\alpha = d + \mu + \frac{c}{m+h}$

They are feasible if $\beta > 4d\alpha^2/A^2$; then

$$I_4 = \frac{A}{2\alpha} - \left(\frac{A^2}{4\alpha^2} - \frac{d}{\beta}\right)^{1/2}, \quad I_5 = \frac{A}{2\alpha} + \left(\frac{A^2}{4\alpha^2} - \frac{d}{\beta}\right)^{1/2}.$$



FIGURE 1. Hopf bifurcation diagram of the stability of equilibria E_4 and E_5 (2.2) of the dynamical system (2.1) in the (β, c) plane, based on the Routh-Hurwitz conditions (2.3) and with (2.4). In region I, the conditions (2.3) for $e = e_i(\beta, c)$, i = 1, 2, 3, are not satisfied, and in region II they are satisfied.

2.2. Behavior of the dynamical system. The point E_1 corresponds to the extinction of the epidemic. On the other hand, it is obtained in [39] that E_2 is unstable and E_3 is stable. To handle the dynamics of the system around E_4 and E_5 , we first recall that at E_5 , the stability matrix is given by

$$\boldsymbol{J}^{*}(E_{5}) = \begin{bmatrix} -\beta I_{5}^{2} - d & -\frac{2c}{m+h} - 2d - 2\mu & 0\\ \beta I_{5}^{2} & \frac{2c}{m+h} + d + \mu - \frac{cm}{(m+h)^{2}} & -\frac{ch^{2}}{(m+h)^{2}}\\ 0 & \delta/h & -\delta \end{bmatrix},$$

and that $J^*(E_4)$ is given by the same expression with I_4 in place of I_5 . The characteristic equation associated with possible equilibria E_4 and E_5 is

$$\det(\boldsymbol{J}^* - \lambda \boldsymbol{I}) = 0 \Leftrightarrow \lambda^3 + e_1 \lambda^2 + e_2 \lambda + e_3 = 0,$$

where $e_1 = -\operatorname{tr}(\boldsymbol{J}^*)$, $e_2 = \sum_{i=1}^3 M_i$, where M_i is the determinant of the matrix obtained by eliminating row *i* and column *i* from \boldsymbol{J}^* , and $e_3 = -\det \boldsymbol{J}^*$. Here, the Routh-Hurwitz conditions (cf., e.g., [17, Sect. 6.4]) mean that there are only roots with strictly negative real parts, and hence the corresponding state is stable, when

$$e_1 > 0, \quad e_3 > 0, \quad \text{and} \quad e_1 e_2 > e_3.$$
 (2.3)

This criterion is applied to $J^*(E_5)$. In order to determine a set of parameters, where scenarios will be studied to determine the effect of considering different values of the infectious force of the epidemic expressed by the parameter β and of the search rate c of the predators towards infected prey, we fix the parameters

$$A = 1, \quad \mu = 1.8, \quad d = 1, \quad m = 30, \quad h = 0.1, \quad \delta = 0.4,$$
 (2.4)

6

where the values of A, μ , and d are those proposed in [39] while the choices of m, h, and δ are assumed but are comparable with those utilized in [20]. To determine suitable choices of β and c, we assume that $e_1 = e_1(\beta, c)$, $e_2 = e_2(\beta, c)$ and $e_3 = e_3(\beta, c)$ and analyze satisfaction of the Routh-Hurwitz condition (2.3). This gives rise to two distinct regions in the (β, c) -plane, see Figure 1. A Hopf bifurcation occurs across the curve that separates these regions.

2.3. **Spatio-temporal predator-prey model.** The ODE model (2.1) exhibits a rich solution behaviour, which motivates us to analyze a spatial variant that in turn generalizes the spatio-temporal hyperbolic-parabolic predator-prey model studied by Colombo and Rossi [13]. Following [12, 13], we then replace (2.1c) by a non-local spatio-temporal PDE that contains a term describing the spatial movement of predators towards infected prey, and equip (2.1a) and (2.1b) with diffusion terms. The resulting spatio-temporal model is given by

$$\frac{\partial S}{\partial t} - D_S \Delta S = A - dS - \beta S I^2, \qquad (2.5a)$$

$$\frac{\partial I}{\partial t} - D_I \Delta I = \beta S I^2 - (d+\mu)I - \frac{cYI}{mY+I},$$
(2.5b)

$$\frac{\partial Y}{\partial t} + \nabla \cdot \boldsymbol{F}_Y \left(S, I, Y \right) = \delta Y \left(1 - \frac{hY}{I} \right), \qquad (2.5c)$$

where $\nabla \cdot$ denotes the (spatial) divergence operator. Here x and y are the space variables, $\Delta = \partial^2/\partial x^2 + \partial^2/\partial y^2$ is the two-dimensional Laplace operator, and D_S and D_I are the diffusion coefficients, i.e. we assume standard linear diffusion for the prey compartments S and I.

The right-hand sides of (2.5) are identical to that of the non-spatial ODE model (2.1), i.e., this model is recovered if all divergence terms on the left-hand sides are set to zero and variables are considered to depend on t only, and the unknowns represent suitably scaled densities.

2.4. Convective fluxes and diffusion matrix. The flux F_Y appearing in the left-hand side of (2.5c) is assumed to have the form $F_Y(I, Y) = \kappa_Y Y V(I)$, where $\kappa_Y \geq 0$ is constant and

$$\boldsymbol{V}(w) = \frac{\nabla(w*\eta)}{\sqrt{1+\|\nabla(w*\eta)\|^2}}$$
(2.6)

is the non-local unscaled velocity function [13]. Here η denotes a radial convolution kernel with radius ε , i.e., η is a piecewise smooth function such that

$$\eta(\boldsymbol{x}) = \eta(\|\boldsymbol{x}\|_2), \quad \eta(\boldsymbol{x}) \ge 0, \quad \eta(\boldsymbol{x}) = 0 \text{ for } \|\boldsymbol{x}\| > \varepsilon, \quad \text{and } \int_{\mathbb{R}^2} \eta(\boldsymbol{x}) \, \mathrm{d}\boldsymbol{x} = 1, \quad (2.7)$$

i.e., for any function w defined on $\Omega \times \mathcal{T}$ and $\boldsymbol{x} \in \Omega$ such that $B_{\varepsilon}(\boldsymbol{x}) := \{ \boldsymbol{y} \in \mathbb{R}^2 : \| \boldsymbol{y} - \boldsymbol{x} \| < \varepsilon \} \subset \Omega$, we have

$$(w(\cdot,t)*\eta)(\boldsymbol{x}) = \int_{B_{\varepsilon}(\boldsymbol{x})} w(\boldsymbol{y},t)\eta(\boldsymbol{x}-\boldsymbol{y}) \,\mathrm{d}\boldsymbol{y} = \int_{\mathbb{R}^2} w(\boldsymbol{y},t)\eta(\boldsymbol{x}-\boldsymbol{y}) \,\mathrm{d}\boldsymbol{y}$$

(with slight modifications for points \boldsymbol{x} with $\operatorname{dist}(\boldsymbol{x}, \partial \Omega) < \varepsilon$.) Since $\nabla(\boldsymbol{w} * \eta) = \boldsymbol{w} * \nabla \eta$, the velocity $\boldsymbol{V}(\boldsymbol{w})$ indeed depends (non-locally) on \boldsymbol{w} and not on its gradient.

Summarizing, we obtain $\mathbf{F}^{c}(\mathbf{u}) = (0, 0, F_{Y}(I, Y))^{T}$ and $\mathcal{D} = \text{diag}(D_{S}, D_{I}, 0)$. The vector $\mathbf{s}(\mathbf{u}) = (s_{1}(\mathbf{u}), s_{2}(\mathbf{u}), s_{3}(\mathbf{u}))^{T}$ is given by the right-hand sides of (2.5). The system (2.5) is considered on $\Omega \times \mathcal{T}$ along with the initial condition

$$\boldsymbol{u}(\boldsymbol{x},0) = \boldsymbol{u}_0(\boldsymbol{x}), \quad \boldsymbol{x} \in \Omega,$$
(2.8)

where u_0 is a given function, and zero-flux boundary conditions

$$(\mathbf{F}^{c}(\mathbf{u}) - \mathbf{\mathcal{D}}\nabla\mathbf{u}) \cdot \mathbf{n} = \mathbf{0}, \quad \mathbf{x} \in \partial\Omega, \quad t \in (0, T],$$
 (2.9)

where \boldsymbol{n} is the unit exterior normal vector to the boundary $\partial \Omega$ of Ω .

3. Numerical method.

3.1. Spatial discretization of local convection and diffusion terms. We take $\Omega = [0, L] \times [0, L]$ and use a Cartesian grid with nodes (x_i, y_j) , $i, j = 1, \ldots, M$, with $x_i = y_i = (i - 1/2)h$, h = L/M. We discretize $\nabla \cdot \mathbf{F}^{c}(\mathbf{u})$ by WENO finite differences and $\Delta \mathbf{u}$ by the standard second-order scheme with a five-point stencil to get a spatial semi-discretization of (1.1) for a $3 \times M \times M$ -matrix $\mathbf{v}(t)$ of unknown approximations $v_{\ell,i,j}(t) \approx u_{\ell}(x_i, y_j, t)$ for $i, j = 1, \ldots, M$ and $\ell = 1, \ldots, 3$. This spatial semi-discretization is given by

$$\boldsymbol{v}' = -\nabla_h \cdot \tilde{\boldsymbol{F}}^{c}(\boldsymbol{v}) + \boldsymbol{\mathcal{B}}\boldsymbol{v} + \boldsymbol{S}(\boldsymbol{v}).$$
(3.1)

Applying a suitable IMEX-RK scheme to (3.1) yields the final fully-discrete scheme (see Section 3.4). Here $\nabla_h \cdot \tilde{\boldsymbol{F}}^c(\boldsymbol{v})$ is the discretization of $\nabla \cdot \boldsymbol{F}^c(\boldsymbol{u})$, to be defined in Section 3.3, and

$$(\mathcal{B}\boldsymbol{v})_{\ell,i,j} = \mu_{\ell}(\boldsymbol{\Delta}_{h}\boldsymbol{v}_{\ell})_{i,j}, \quad i,j = 1,\dots, M, \quad \ell = 1,\dots, 3$$
(3.2)

denotes the discretization of the diffusion terms. Here v_{ℓ} denotes the $M \times M$ submatrix given by $(v_{\ell})_{i,j} = v_{\ell,i,j}$ and Δ_h is the approximation of the standard twodimensional Laplacian operator with Neumann boundary conditions. Furthermore, S(v) is the $3 \times M \times M$ -matrix with components

$$S(v)_{\ell,i,j} = s_{\ell}(v_{\ell,i,j}), \quad i, j = 1, \dots, M, \quad \ell = 1, \dots, 3,$$

with corresponding submatrices $S_{\ell}(v)$, given by $S_{\ell}(v)_{i,j} = s_{\ell}(v_{\ell,i,j})$.

3.2. Discretization of the convolutions. The following identity arises from (2.6) if we take into account that $\nabla(w * \eta) = w * \nabla \eta$:

$$\boldsymbol{V}(w) = \frac{w * \boldsymbol{\nu}}{\sqrt{1 + \|w * \boldsymbol{\nu}\|^2}}, \quad \boldsymbol{\nu} = \left(\frac{\partial \eta}{\partial x}, \frac{\partial \eta}{\partial y}\right). \tag{3.3}$$

The corresponding convolutions $w * \partial \eta / \partial x$ and $w * \partial \eta / \partial y$ are calculated approximately on the discrete grid via a composite Newton-Cotes quadrature formula, such as the composite Simpson rule.

Since $B_{\varepsilon}(0) \subseteq [-rh, rh]^2$, $r = \lceil \varepsilon/h \rceil < M$, and according to boundary conditions, w is extended to the exterior of Ω by reflection, e.g. by setting w(-x, y) = w(x, y), $(x, y) \in \Omega$, we obtain

$$(w * \chi)(x_i, y_j) \approx \sum_{p=-r}^r \sum_{q=-r}^r \beta_{p,q} w(x_{i-p}, y_{j-q}), \quad \beta_{p,q} = h^2 \alpha_p \alpha_q \chi(x_p, y_q), \quad (3.4)$$

where α_p and α_q are the coefficients in the quadrature rule (e.g., for the composite Simpson rule, $\boldsymbol{\alpha} = (1, 4, 2, 4, \dots, 2, 4, 1)$). Consequently, the approximation (3.4) for $\boldsymbol{W} = (w_{i,j}) \in \mathbb{R}^{M \times M}$, $w_{i,j} \approx w(x_i, y_j)$, is given by

$$(w * \chi)(x_i, y_j) \approx (\mathbf{W} *_h \beta)_{i,j} := \sum_{p=-r}^r \sum_{q=-r}^r \beta_{p,q} w_{[i-p]_M, [j-q]_M},$$
 (3.5)

where we define

$$[i]_M := \begin{cases} -i+1 & \text{for } -r+1 \le i \le 0, \\ i & \text{for } 1 \le i \le M, \\ 2M+1-i & \text{for } M+1 \le i \le M+r. \end{cases}$$

The discrete approximation of V(w) in (3.3) is then given by

$$oldsymbol{V}_h(oldsymbol{W}) = rac{oldsymbol{W}*_holdsymbol{
u}}{\sqrt{1+\|oldsymbol{W}*_h
u\|^2}}$$

Since $r \approx \varepsilon/h = \varepsilon M/L$, the computational cost of this discrete convolution is $M^2(2r+1)^2 \approx 4\varepsilon^2 M^4/L^2$, which can be very high for large M. This cost can be substantially reduced to $\mathcal{O}(M^2 \log M)$ by performing a convolution with periodic data by Fast Fourier Transforms (FFTs) (see [23, 37]). To achieve this goal, we define from $\mathbf{W} = (w_{i,j}) \in \mathbb{R}^{M \times M}$ a matrix $\tilde{\mathbf{W}} = (\tilde{w}_{i,j}) \in \mathbb{R}^{2M \times 2M}$ such that

$$\tilde{w}_{i,j} = w_{[i]_M,[j]_M}, \quad i, j = 1, \dots, 2M$$

and use the notation $[i]'_{2M} = \mod(i-1, 2M) + 1$, i.e., $[i]'_{2M} = i + 2kM$, with k being the integer such that $1 \leq [i]'_{2M} \leq 2M$. It is then readily checked that

$$w_{[i]_M,[j]_M} = \tilde{w}_{[i]'_{2M},[j]'_{2M}}, \quad i,j = -r+1,\dots,M+r.$$

Therefore (3.5) for i, j = 1, ..., M can be rewritten as

$$(\boldsymbol{W} *_{h} \beta)_{i,j} = \sum_{p=-r}^{r} \sum_{q=-r}^{r} \beta_{p,q} \tilde{w}_{[i-p]'_{2M},[j-q]'_{2M}}.$$
(3.6)

The convolution on the right-hand side of (3.6) can be performed by FFTs applied to the $(2M) \times (2M)$ matrix $\tilde{\boldsymbol{W}}$. To save further computational costs, the FFT of the kernel $\beta_{p,q}$ is performed only once, son each convolution entails two twodimensional FFT of $(2M) \times (2M)$ matrices and a product of $4M^2$ numbers, with an overall computational cost of $\mathcal{O}(M^2 \log M)$.

3.3. Discretization of the convective term. The convective flux for the prey compartments (S and I, corresponding to $\ell \in \{1,2\}$) is zero and for the predator (species Y, $\ell = 3$) is given by $\mathbf{F}_3^c(\mathbf{u}) = u_3(\kappa_3 \mathbf{V}(u_2))$. To discretize its divergence $\nabla \cdot \mathbf{F}_3^c(\mathbf{u})$ for the approximation \mathbf{v} , we first approximate the convolution terms as expounded in Section 3.2 to obtain

$$\tilde{\boldsymbol{F}}_{3}^{c}(\boldsymbol{v})_{i,j} = \boldsymbol{v}_{\ell,i,j} \left(\kappa_{3} \boldsymbol{V}_{h} \left(v_{2} \right)_{i,j} \right) \in \mathbb{R}^{2}.$$

We introduce the notation $(f_{i,j}^x, f_{i,j}^y) := \tilde{F}_{\ell}^c(\boldsymbol{v})_{i,j}$, where we have dropped the index ℓ in $f_{i,j}^x$ etc. to obtain clearer expressions. Our purpose is to use a fifth-order WENO finite difference discretization [22, 29, 36] of $\nabla \cdot F_{\ell}^c(\boldsymbol{u})$ for which

$$\nabla \cdot \boldsymbol{F}_{\ell}^{c}(\boldsymbol{u})(x_{i}, y_{j}) \approx \nabla_{h} \cdot \tilde{\boldsymbol{F}}^{c}(\boldsymbol{v})_{\ell, i, j} := \frac{\hat{f}_{i+1/2, j}^{x} - \hat{f}_{i-1/2, j}^{x}}{h} + \frac{\hat{f}_{i, j+1/2}^{y} - \hat{f}_{i, j-1/2}^{y}}{h}$$

for suitable numerical fluxes $\hat{f}_{i+1/2,j}^x$, $\hat{f}_{i,j+1/2}^y$ obtained by WENO reconstructions of split fluxes. For the numerical flux in the *x*-direction, the Lax-Friedrichs-type flux splitting $f^{x,\pm}$ is given by

$$f_{i,j}^{x,\pm} = \frac{1}{2} \left(f_{i,j}^x \pm \alpha^x v_{\ell,i,j} \right), \quad \alpha^x = \max_{i,j} \left| \boldsymbol{V}_h^x(\boldsymbol{v})_{i,j} \right|,$$

with an analogous formula for $f_{i,j}^{y,\pm}$. If \mathcal{R}^{\pm} denotes fifth-order WENO upwind biased reconstructions and we use matlab-type notation for submatrices, then

$$\begin{split} \hat{f}^x_{i+1/2,j} &= \mathcal{R}^+ \big(f^{x,+}_{i-2:i+2,j} \big) + \mathcal{R}^- \big(f^{x,-}_{i-1:i+3,j} \big), \\ \hat{f}^y_{i,j+1/2} &= \mathcal{R}^+ \big(f^{y,+}_{i,j-2:j+2} \big) + \mathcal{R}^- \big(f^{y,-}_{i,j-1:j+3} \big). \end{split}$$

3.4. Implicit-explicit Runge-Kutta schemes. To specify the IMEX-RK integrators for (3.1), we rewrite (3.1) as (1.2), where

$$\Phi^*(\boldsymbol{v}) := -\nabla_h \cdot \tilde{\boldsymbol{F}}^{c}(\boldsymbol{v}) + \boldsymbol{S}(\boldsymbol{v}), \qquad \Phi(\boldsymbol{v}) := \boldsymbol{\mathcal{B}}\boldsymbol{v}.$$
(3.7)

The diffusive part $\Phi(\boldsymbol{v})$ is handled by an implicit *s*-stage diagonally implicit (DIRK) scheme with coefficients $\boldsymbol{A} \in \mathbb{R}^{s \times s}$, $\boldsymbol{b}, \boldsymbol{c} \in \mathbb{R}^{s}$, in the common Butcher notation, where $\boldsymbol{A} = (a_{ij})$ with $a_{ij} = 0$ for j > i. For the term $\Phi^*(\boldsymbol{v})$ we employ an *s*-stage explicit scheme with coefficients $\hat{\boldsymbol{A}} \in \mathbb{R}^{s \times s}$, $\hat{\boldsymbol{b}}, \hat{\boldsymbol{c}} \in \mathbb{R}^{s}$ and $\hat{\boldsymbol{A}} = (\hat{a}_{ij})$ with $\hat{a}_{ij} = 0$ for $j \geq i$. In our simulations, we limit ourselves to the second-order IMEX-RK scheme H-DIRK2(2,2,2) that corresponds to the pair of Butcher arrays

$$\begin{array}{c|c|c|c} \hline c & A \\ \hline & 1/2 & 1/2 & 0 \\ \hline & b^{\mathrm{T}} \end{array} = \begin{array}{c|c|c|c} 1/2 & 1/2 & 0 & & \\ \hline 1/2 & 0 & 1/2 & & \\ \hline & 1/2 & 1/2 \end{array} , \qquad \begin{array}{c|c|c|c|c|c} \hline c & \hat{A} & & \\ \hline & \hat{b}^{\mathrm{T}} \end{array} = \begin{array}{c|c|c|c} 0 & 0 & 0 & & \\ \hline 1 & 1 & 0 & & \\ \hline & 1/2 & 1/2 \end{array}$$

Alternative choices are provided and discussed in [5,6,33]. If applied to the equation (1.2), the IMEX-RK scheme gives rise to the following algorithm (see [33]).

Algorithm 3.1.

10

Input: approximate solution vector \mathbf{v}^n of (1.2) for $t = t^n$ for p = 1, ..., sif p = 1 then $\hat{\mathbf{v}}^{(1)} \leftarrow \mathbf{v}^n$, $\bar{\mathbf{v}}^{(1)} \leftarrow \mathbf{v}^n$ else compute $\hat{\mathbf{v}}^{(p)}$ and $\bar{\mathbf{v}}^{(p)}$ from the known values $\mathbf{K}_1, ..., \mathbf{K}_{p-1}$: $\hat{\mathbf{v}}^{(p)} \leftarrow \mathbf{v}^n + \Delta t \sum_{j=1}^{p-1} \hat{a}_{pj} \mathbf{K}_j$, $\bar{\mathbf{v}}^{(p)} \leftarrow \mathbf{v}^n + \Delta t \sum_{j=1}^{p-1} a_{pj} \mathbf{K}_j$

endif

solve for \mathbf{K}_p the linear system

$$\boldsymbol{K}_{p} = \Phi^{*}(\boldsymbol{\hat{v}}^{(p)}) + \Phi(\boldsymbol{\bar{v}}^{(p)} + \Delta t a_{pp} \boldsymbol{K}_{p})$$
(3.8)

endfor

$$\boldsymbol{v}^{n+1} \leftarrow \boldsymbol{v}^n + \Delta t \sum_{j=1}^s b_j \boldsymbol{K}_j$$

Output: approximate solution vector v^{n+1} of (1.2) for $t = t^{n+1} = t^n + \Delta t$.

To solve the linear equation (3.8) for K_p , in view of (3.7) we rewrite it as

$$(\boldsymbol{I} - \Delta t a_{pp} \boldsymbol{\mathcal{B}}) \boldsymbol{K}_{p} = \boldsymbol{b}^{(p)}, \quad \boldsymbol{b}^{(p)} := \Phi^{*} \left(\hat{\boldsymbol{v}}^{(p)} \right) + \boldsymbol{\mathcal{B}} \bar{\boldsymbol{v}}^{(p)}, \tag{3.9}$$

where I denotes the identity operator for $3 \times M \times M$ matrices. From the definition of the matrix \mathcal{B} in (3.2) and from the definition of Φ^* in (3.7), if we equate the ℓ submatrices along the first dimension of both sides of (3.9) we get

$$(\boldsymbol{I}_{M \times M} - \Delta t a_{pp} \mu_{\ell} \boldsymbol{\Delta}_{h}) (\boldsymbol{K}_{p})_{\ell} = -\nabla_{h} \cdot \tilde{\boldsymbol{F}}_{\ell}^{c} (\hat{\boldsymbol{v}}^{(p)}) + \boldsymbol{S}_{\ell} (\hat{\boldsymbol{v}}^{(p)}) + \mu_{\ell} \boldsymbol{\Delta}_{h} \bar{\boldsymbol{v}}_{\ell}^{(p)}, \quad \ell = 1, 2, 3,$$

$$(3.10)$$



FIGURE 2. Numerical solution of the ODE model (2.1) with parameters (2.4) and (4.1), starting from S(0) = 0.2, I(0) = 0.3, and Y(0) = 0.3. The blue curve in the right plot is the solution curve in (S, I, Y)-space. The other curves represent projections into the (S, I)-, (Y, S)- and (Y, I)-planes.

where $I_{M \times M}$ is the $M \times M$ identity operator and, e.g., $(\mathbf{K}_p)_{\ell}$ is the ℓ submatrix of \mathbf{K}_p along the first dimension, i.e., $((\mathbf{K}_p)_{\ell})_{i,j} = (\mathbf{K}_p)_{\ell,i,j}$. If $\mu_{\ell} = 0$ (for $\ell = 3$), then

$$(oldsymbol{K}_p)_\ell = -ig(
abla_h \cdot ilde{oldsymbol{F}}^{ ext{c}}(oldsymbol{\hat{v}}^{(p)}) ig)_\ell + oldsymbol{S}_\ell(oldsymbol{v}),$$

otherwise (3.10) is solved by Fast Cosine Transforms (due to boundary conditions), which entails a nearly optimal computational cost of $\mathcal{O}(M^2 \log M)$.

4. Numerical results. We wish to compare the ODE model (2.1) with the spatiotemporal model (2.5) in different scenarios.

4.1. Solution to the ODE model. For this model we consider parameter values (2.4) along with

$$\beta = 42, \quad c = 5.$$
 (4.1)

This choice is in accordance with the region of stability shown in Figure 1 (these parameters will be varied within the numerical solution of spatio-temporal models). For the parameters (2.4) and (4.1), we obtain the following equilibria of Theorem 2.1 (besides $E_1 = (1, 0, 0)$) where the predator is absent:

$$E_2 = (0.7516611, 0.0886924, 0), \quad E_3 = (0.2483389, 0.2684504, 0), \quad (4.2)$$

as well as the two points

 $E_4 = (0.7013155, 0.1006989, 1.0069894), \quad E_5 = (0.2986844, 0.2364426, 2.3644263).$ Since all eigenvalues of

$$\boldsymbol{J}^{*}(E_{5}) = \begin{bmatrix} -3.3480149 & -5.9322259 & 0\\ 2.3480149 & 2.9666648 & -0.000055\\ 0 & 4 & -0.4 \end{bmatrix}$$

12



FIGURE 3. Examples 1 to 10: initial data $S_0(x, y)$, $I_0(x, y)$, and $Y_0(x, y)$ (a) for Examples 1, 4, and 7, (b) for Examples 2, 5, and 8, (c) for Examples 3, 6, 9, and 10.

have negative real part, the equilibrium E_5 is stable. On the other hand, $J^*(E_4)$ has an eigenvalue with positive real part and therefore the point E_4 is unstable. As a consequence, whenever predators and prey are present the solution orbits of (2.1) approach the stable equilibrium E_5 , as is illustrated in Figure 2.

4.2. Solution to the spatio-temporal model. In Examples 1 to 10 we select the spatial domain $\Omega = [0, L] \times [0, L]$ with L = 100. For the spatial discretization we use M = 400, such that $\Delta x = L/M = 0.25$. For the time discretization we employ Algorithm 3.1. The parameters $\varepsilon = 4$ and $\kappa_Y = 1$ are chosen in the nonlocal term, and the diffusion constants are chosen as $D_S = 6$ and $D_I = 1$, corresponding to the assumption that infected individuals of the prey population exhibit a lower degree of mobility than their susceptible counterparts. The initial condition for S and I is always a spatially distributed random perturbation of the respective values 0.3 and 0.2. Three cases will be considered for the initial condition for predators, namely (a) absence of predators, (b) a "triangular" initial condition for predators to describe how predators invade a region initially occupied by prey only, and (c) a random initial condition for predators. These three scenarios are illustrated in Figure 3. The simulations are carried out until the nonlocal system (2.5) attains a stable non-homogeneous steady state.

Furthermore, we wish to compare the numerical results with the predictions made by the non-spatial ODE model. To this end we determine for each compartment



FIGURE 4. Example 1: numerical solution of (2.5) in absence of predators with parameters (2.4) and (4.1) at three different times. The initial datum is given in Figure 3 (a).

 $X \in \{S, I, Y\}$ and time instants $t^n = n\Delta t$ the following quantity:

$$\mathcal{I}(X) = \mathcal{I}(X, t^n) := h^2 \sum_{i,j=1}^{N} X_{i,j}^n \approx \int_{\Omega} X(\boldsymbol{x}, t^n) \,\mathrm{d}\boldsymbol{x}, \tag{4.3}$$

which represents the approximate total number in Ω of individuals of compartment X at time t^n . We recall that in the PDE model, the unknowns $X \in \mathcal{C}$ are *densities*, and so an integration over Ω is necessary to make results comparable with those of a model that predicts the *total number* of individuals in each compartment (as does the dynamical system (2.1)).

Example 1. Here, we consider absence of the predators, i.e., only the equations (2.5a), (2.5b) are considered, with the parameters given by (2.4) and (4.1). The values of β and μ in combination with $D_S = 6$ and $D_I = 1$ have been chosen such that they lie in the Turing region, see [39, Example 6], which means that the formation of a permanent spatial pattern by the standard Turing mechanism in reaction-diffusion systems (see, e.g., [31]) is expected. In our simulation the initial conditions for S and I are those of Figure 3 (a). In Figure 4 we display the numerical solution at three different times. We observe the formation of a pattern of spots similar to that observed in [39]. This distribution will remain constant over time. Moreover, Figure 5 shows the integral quantities $\mathcal{I}(S)$, $\mathcal{I}(I)$, and wherever appropriate $\mathcal{I}(Y)$, for Examples 1 to 6. For Example 1, and consistently with the nearly stationary spatial pattern, we observe (in Figures 5 (a) and (b)) that $\mathcal{I}(S)$ and $\mathcal{I}(I)$ become practically stationary after very short time, and assume values close to $L^2 \times 0.2483389 \approx 2483$ and $L^2 \times 0.2684504 \approx 2684$ that would be

14



FIGURE 5. Examples 1 to 6: integral quantities (4.3).

obtained if the solution were equal to the stable equilibrium E_3 (see (4.2)) on the whole domain. In this sense, in this example the global behaviour of the model is consistent with (2.1).



FIGURE 6. Example 2: numerical solution of (2.5) with parameters (2.4) and (4.1) at three different times. The initial datum is given in Figure 3 (b).

Example 2. Here we introduce predators through the "triangular" initial condition of Figure 3 (b). All other parameters are the same as in Example 1. In Figure 6 we display the dynamics of model (2.5). The convective term in (2.5c) defined in terms of the non-local velocity function (2.6) allows the movement of the predators to places where infective prey are present. At simulated time t = 30 we observe that the distribution of prey varies in comparison with Example 1 in places where predators are present. At t = 90 we observe that in places where the predators are absent, preys are distributed in forming spots as in Example 1, however, where predators are present, these patterns take the form of stripes. At t = 200, predators are present in the whole domain, and this has motivated that the distribution of the prey in the domain varies drastically from its natural distribution without predator (Example 1). This distribution will remain constant over time. The corresponding plots of $\mathcal{I}(S)$ and $\mathcal{I}(I)$ (Figure 5 (c)) and $\mathcal{I}(Y)$ indicate a growth of the predator population over a very long period of time. Shortly before t = 200 these quantities become constant. It is worth noting that while $\mathcal{I}(S)$ and $\mathcal{I}(I)$ tend to values that



FIGURE 7. Example 3: numerical solution of (2.5) with parameters (2.4) and (4.1) at three different times. The initial datum is given in Figure 3 (c).

are close to L^2 times the first and second component of E_5 , namely 2986.8 and 2364.4, respectively, the limit value of $\mathcal{I}(Y)$ exceeds L^2 times the Y-component of E_5 , that is 23644.3, by more than a factor of two.

Example 3. Here we consider a random initial condition for predators with the same parameters of Example 1. The initial condition is shown in Figure 3 (c). In Figure 7 we display the dynamics of the model with this initial condition. At simulated time t = 30 we observe that stripes and some spots patterns emerge mixed in the distribution of each species. After the stripe patterns form, they grow steadily with time until they reach certain arm length, and the spatial patterns become distinct at t = 200. It is observed that these stripe patterns are not similar to those obtained in Example 1, instead these patterns in form of stripes or filaments are very similar to those observed in Example 2. Furthermore, Figures 5 (e) and (f) indicate that $\mathcal{I}(S)$, $\mathcal{I}(Y)$ and $\mathcal{I}(Y)$ quickly attain steady states, with final values similar to those of Example 2.

PREDATOR-PREY MODEL WITH INFECTED PREY



FIGURE 8. Example 4: numerical solution of (2.5) with parameters (2.4) and (4.4) at three different times. The initial datum is given in Figure 3 (a).

Example 4. Here, we consider absence of predators like in Example 1 and we use here and in Examples 5 and 6 the parameter values

$$\beta = 46, \quad c = 4.$$
 (4.4)

In this case the parameters β and μ together with the diffusion constants are not in the Turing space. The steady states according to Theorem 2.1 are now, besides $E_1 = (1, 0, 0)$,

$$E_{2} = (0.7820731, 0.0778310, 0), \quad E_{3} = (0.2179270, 0.2793118, 0), \quad (4.5)$$

$$E_{4} = (0.7510048, 0.0848976, 0.8489759), \quad E_{5} = (0.2489953, 0.2560630, 2.5606299). \quad (4.6)$$

Considering the initial conditions of Figure 3 (a), we obtain in this case that no patterns are formed and the system quickly arrives at a constant equilibrium (uniform distribution for S and I in whole domain), which is maintained over time, as can be observed in Figure 8. This figure, as well as Figures 5 (g) and (h), indicate that the global equilibrium is E_3 given by (4.5).

Example 5. In this case we consider the same scenario as in Example 2, namely the triangular initial condition of Figure 3 (b), but utilize the parameters (4.4) instead of (4.1). In this case, as in Example 2, predators are heading towards the infected prey, and as is shown in Figure 9, the whole domain becomes successively filled with a pattern formed of stripes. Note that in regions occupied by the prey before the arrival of predators the solution is constant (see Example 4), and in particular does not exhibit any formed pattern (in contrast to Example 2). In a sense this scenario illustrates how the model (2.5), under the appropriate choice of parameters, predicts



FIGURE 9. Example 5: numerical solution of (2.5) with parameters (2.4) and (4.4) at three different times. The initial datum is given in Figure 3 (b).

the formation of spatial patterns among the prey upon arrival of the predators. Furthermore, we observe that a "front" of predators is moving into the domain initially occupied by prey only, which seems to move at a lower velocity than in Example 2. It is worth noting that the final distribution of prey corresponds to total S and I populations close to L^2 times the corresponding entries of E_5 given by (4.6), while as in previous cases the final population of predators is much higher than L^2 times the Y-component of E_5 .

Example 6. This case is the analogue of Example 2, namely we impose the initial condition of Figure 3 (c). In Figure 10 we display the dynamics of the model with this initial condition. We observe that at simulated time t = 50, some stripes and spots emerge as patterns in the distribution of each species. Comparing the successive solutions for t = 50, t = 100 and t = 200, we observe that once the stripes form, they grow steadily in time incorporating "spots", until they reach certain arm length, and the spatial patterns become slightly distinct at t = 200 since at this time there are strips and spots. It is observed that these patterns differ

PREDATOR-PREY MODEL WITH INFECTED PREY



FIGURE 10. Example 6: numerical solution of (2.5) with parameters (2.4) and (4.4) at three different times. The initial datum is given in Figure 3 (c).

from those obtained in Example 5, where only stripe-like patterns are formed. The observations corresponding to the evolution of $\mathcal{I}(S)$, $\mathcal{I}(I)$ and $\mathcal{I}(Y)$ are analogous to those of Example 3.

Example 7. Here and in Examples 8 and 9 we use the values

$$\beta = 46, \quad c = 10.$$
 (4.7)

The equilibrium points are $E_1 = (1, 0, 0)$, E_2 and E_3 given by (4.5), and

 $E_4 = (0.6916269, 0.984517, 0.9845173), \quad E_5 = (0.3083731, 0.2208100, 2.2081005).$

The integral quantities for this case and Examples 8 to 10 are plotted in Figure 12.

As in Example 4, the parameters β and μ together with the diffusion constants are not in the Turing space. Considering the initial condition of Figure 3 (a), it is obtained that in this case no patterns are formed and the system rapidly tends to constant equilibrium which is maintained in time, see Figure 11. For this case we observe a behavior similar to that of Example 7.



FIGURE 11. Example 7: numerical solution of (2.5) with parameters (2.4) and (4.7) at three different times. The initial datum is given in Figure 3 (a).

Example 8. This case is based on the same initial "triangular" scenario (Figure 3 (b)) as Examples 2 and 5. The numerical solution (Figure 13) shows that the predators are heading towards the infected prey, and for t = 50 and t = 150 we observe that the distribution of the prey varies in comparison with the uniform distribution of Example 7 in those places were the predators are present. The final spatial configuration is marked by spots and stripes for the prey as well as for the predators, and remains in time. However, the stucture of patterns that are formed differs from the results of Example 5. It is interesting to note that the stripes visible at t = 600 are still roughly aligned with the original "front" of predators separating the triangular region from the rest of the domain. The conclusions concerning the evolution of $\mathcal{I}(S)$, $\mathcal{I}(I)$ and $\mathcal{I}(Y)$ (see Figure 12 (c) and (d)) are similar to those for Example 5.

Example 9. We now consider an analogue of Examples 3 and 6 corresponding to the initial configuration of Figure 3 (c). In Figure 14 we display the dynamics of the model with this initial condition. At t = 50 we observe that some stripes and spots patterns emerge mixed in the distribution of each species. The pattern distribution is maintained over time after its formation. It is observed that these patterns differ from those obtained in Examples 5, 6, and 8. As in Examples 3 and 6, the quantities $\mathcal{I}(S), \mathcal{I}(I)$ and $\mathcal{I}(Y)$ very quickly attain constant values (see Figures 12 (e) and (f)).

Example 10. In this example, a constant value is considered for $\beta = 48$ and in combination with the three different values c = 4, c = 6 and c = 10. For these three pairs that are in the region of stability shown in the Figure 1 a variation is obtained in the formation of patterns for the different values of c. The initial datum is given

20



FIGURE 12. Examples 7 to 10: integral quantities (4.3).

by Figure 1(c). In Figure 15 the corresponding numerical results are shown. It turns out that with increasing c the species tend to form more filament-like rather than spot-like spatial structures. Moreover, there is a marked difference in behavior of $\mathcal{I}(S)$, $\mathcal{I}(I)$ and $\mathcal{I}(Y)$, as can be inferred from Figures 12 (g) to (l). Note that



FIGURE 13. Example 8: numerical solution of (2.5) with parameters (2.4) and (4.7) at three different times. The initial datum is given in Figure 3 (b).

the steady-state value of $\mathcal{I}(I)$ decreases with increasing c, as does the steady-state value of $\mathcal{I}(Y)$.

4.3. Convergence test and effect of the choice of ε . In Example 11, we investigate the sensitivity of (2.5) to the variation of discretization parameter $\Delta x = L/M$. We take L = 25 and the model parameters (2.4), (4.4) as in Examples 4 to 6. The initial datum is a random distribution (similar to Figure 1 (c)) defined for a 50 × 50 discretization, which is also utilized for finer discretizations with M = 100, 200, and 400, so the initial condition is exactly the same in all cases. We compute the solution at time t = 15 until we obtain a steady state. The results are displayed in Figure 16. It is observed that the resolution in the solution is improved as Δx is reduced, and that the steady-state solution is apparently the same (modulo, of course, sharpness of resolution) for all values of M. These results do not provide a rigorous convergence proof but indicate that the numerical simulator is reliable.

Finally, in Example 12 we utilize the same scenario as in Example 11, and now fix the spatial discretization M = 400 but vary the parameter ε , that is the radius of the



FIGURE 14. Example 9: numerical solution of (2.5) with parameters (2.4) and (4.7) at three different times. The initial datum is given in Figure 3 (c).

convolution kernel of the nonlocal velocity function (see (2.7)). The corresponding results are given in Figures 17 and 18. We observe that the results for the prey compartments S and I are practically the same for all values of ε condered, while the prey distribution depends appreciably on ε . As ε decreases, the regions in which predators concentrate become smaller and the densities become higher. Moreover, Figures 18 (b), (d), (f), and (h) indicate a decrease of the steady state value of \mathcal{I} as ε is decreased.

5. Conclusions. We have shown that a relatively simple three-species model can produce complex spatio-temporal patterns of predators and prey. As is stated in Section 1.1, the treatment combines the spatio-temporal epidemic model of [39], the non-spatial three-species predator-prey model analyzed in [20], the non-local velocity function introduced in [13] and the numerical method developed in [12]. The numerical method is not fully analyzed herein, and certainly leaves potential for further improvements. For instance, its accuracy deserves further investigation (particularly to establish that its order of convergence corresponds to its design



FIGURE 15. Example 10: variation of the parameter c for $\beta = 48$, showing the numerical solution atr t = 50 for (left) c = 4, (middle) c = 6 and (right) c = 10. All the remaining parameters are as in (2.4). The initial datum is as shown in Figure 3 (c).

order), and we use a second-order spatial discretization of the Laplacian and a second-order time-stepping, which limit the high-order accuracy of the fifth-order WENO spatial semidiscretization. Nevertheless, on the basis of our (limited) evidence of robustness of the scheme (provided by Examples 11 and 12 along with the fact that numerical solutions of Examples 1, 4 and 7 are consistent with those of [39], where, however, the domain Ω has slightly different dimensions), the numerical results allow us to draw some conclusions and conjectures about solutions of (2.5). Let us first point out that long-time stable patterned configurations obtained in our numerical experiments are formed by stripes. This contrasts with the results in [13] obtained for a predator-prey model with one single prey species only, where the system tends to an asymptotic state with regular arrangements of spots of high predator concentrations, in agreement with the radial structure introduced by the nonlocal velocity function and where the diffusion caused by the Laplacian in the prey equation counterbalances the first order nonlocal differential operator in the

24

PREDATOR-PREY MODEL WITH INFECTED PREY



FIGURE 16. Example 11: variation of the spatial discretization $\Delta x = L/M$ of the numerical solution at t = 15 on a domain of side length L = 25.

predator equation [13]. A similar tendency was observed in some scenarios of the eight-compartment epidemiological model studied in [12], where male individuals are supposed to move non-locally in response to the sampled density of their female counterparts. Furthermore, it is calling to attention that in our Example 12, the dominant mechanism of pattern formation seems to be the interaction between susceptible and infected prey, since the solution (see Figure 17) for the prey compartments S and Y is practically independent of ε . That the effect of predator location on the formation of the prey pattern should be weak, at least for the parameters (2.4) and (4.4), can probably be explained by the relative smallness of the predation term on the right-hand side of (2.5b). Finally, it would be interesting to investigate whether the invasion of predators into a domain initially occupied by prey solely, as in Examples 2, 5 and 8, can possibly be modeled by simpler, spatially one-dimensional version of (2.5). Clearly, a more in-depth mathematical analysis to describe the mechanisms that compel these and other phenomena is necessary.

With respect to further extensions, we mention that the spatio-temporal approach could also be applied to the model by Greenhalgh et al. [19] that generalizes (2.1) and [20] in the sense that the assumption that the predator eats infected prey only is replaced by the assumption that the susceptible and infected populations are both exposed to the predator but to varying degrees. Concerning the numerical scheme, further analysis could compare results with higher-order discretizations of the diffusion operator and time-stepping scheme.



FIGURE 17. Example 12: effect of the variation of convolution radius ε on the numerical solution at t = 15 on a domain of side length L = 25.

Acknowledgements. RB is supported by Fondecyt project 1170473; and Centro CRHIAM Proyecto Conicyt Fondap 15130015. PM is supported by Spanish MINECO project MTM2017-83942-P and Conicyt (Chile), project PAI-MEC, folio 80150006. LMV is supported by Fondecyt project 1181511. RB and LMV are also supported by BASAL project CMM, Universidad de Chile and Centro de Investigación en Ingeniería Matemática (CI²MA), Universidad de Concepción and the INRIA Associated Team "Efficient numerical schemes for non-local transport phenomena (NOLOCO; 2018–2020)". EG is supported by CONICYT scholarship. GC acknowledges financial support from grants NSF-IIS RAPID award #1518939, NSF grant 1318788 III: Small: Data Management for Real-Time Data Driven Epidemic simulation, and Conicyt (Chile), project MEC80170119.

REFERENCES

- L.J.S. Allen, B.M. Bolker, Y. Lou, and A.L. Nevai, Asymptotic profiles of the steady states for an SIS epidemic patch model, SIAM J. Appl. Math., 67 (2007), 1283–1309.
- [2] J. Arino, Diseases in metapopulations. In Z. Ma, Y. Zhou, and J. Wu (Eds.), Modeling and Dynamics of Infectious Diseases, Higher Education Press, Beijing, 2009, 64–122.
- [3] J. Arino, J.R. Davis, D. Hartley, R. Jordan, J.M. Miller, and P. van den Driessche, A multispecies epidemic model with spatial dynamics, *Mathematical Medicine and Biology*, 22 (2) (2005), 129–142.
- [4] U. Ascher, S. Ruuth, and J. Spiteri, Implicit-explicit Runge-Kutta methods for time dependent partial differential equations, Appl. Numer. Math., 25 (1997), pp. 151–167.

R. BÜRGER, G. CHOWELL, E. GAVILÁN, P. MULET AND L.M. VILLADA

26



FIGURE 18. Example 12: integral quantities (4.3).

- [5] S. Boscarino, R. Bürger, P. Mulet, G. Russo, and L.M. Villada, Linearly implicit IMEX Runge-Kutta methods for a class of degenerate convection-diffusion problems, *SIAM J. Sci. Comput.*, **37** (2015), B305–B331.
- [6] S. Boscarino, F. Filbet, and G. Russo, High order semi-implicit schemes for time dependent partial differential equations, J. Sci. Comput., 68 (2016), 975–1001.
- [7] S. Boscarino, P.G. LeFloch, and G. Russo, High-order asymptotic-preserving methods for fully nonlinear relaxation problems, SIAM J. Sci. Comput., 36 (2014), A377–A395.
- [8] S. Boscarino and G. Russo, On a class of uniformly accurate IMEX Runge-Kutta schemes and applications to hyperbolic systems with relaxation, SIAM J. Sci. Comput., 31 (2009), 1926–1945.
- [9] S. Boscarino and G. Russo, Flux-explicit IMEX Runge-Kutta schemes for hyperbolic to parabolic relaxation problems, SIAM J. Numer. Anal., 51 (2013), 163–190.
- [10] F. Brauer and C. Kribs, Dynamical Systems for Biological Modeling: An Introduction, CRC Press, Boca Raton, FL, USA, 2016.
- [11] R. Bürger, G. Chowell, P. Mulet, and L.M. Villada, Modelling the spatial-temporal progression of the 2009 A/H1N1 influenza pandemic in Chile, *Math. Biosci. Eng.*, 13 (2016), 43–65.
- [12] R. Bürger, G. Chowell, E. Gavilán, P. Mulet, and L.M. Villada, Numerical solution of a spatio-temporal gender-structured model for hantavirus infection in rodents, *Math. Biosci. Eng.*, **15** (2018), 95–123.
- [13] R.M. Colombo and E. Rossi, Hyperbolic predators versus parabolic preys, Commun. Math. Sci., 13 (2015), 369–400.

- [14] M. Crouzeix, Une méthode multipas implicite-explicite pour l'approximation des équations d'évolution paraboliques, Numer. Math., 35 (1980), 257–276.
- [15] O. Diekmann, H. Heesterbeek, and T. Britton, Mathematical Tools for Understanding Infectious Disease Dynamics, Princeton Series in Theoretical and Computational Biology, Princeton University Press, 2012.
- [16] R. Donat and I. Higueras, On stability issues for IMEX schemes applied to 1D scalar hyperbolic equations with stiff reaction terms, *Math. Comp.*, 80 (2011), 2097–2126.
- [17] L. Edelstein-Keshet, Mathematical Models in Biology, reprint, SIAM, 2005.
- [18] I.M. Foppa, A Historical Introduction to Mathematical Modeling of Infectious Diseases, Academic Press, London, UK, 2016.
- [19] D. Greenhalgh, Q.J.A. Khan, and J.S. Pettigrew, An eco-epidemiological predator-prey modelmwhere predators distinguish between susceptible and infected prey, *Math. Meth. Appl. Sci.*, 40 (2017), 146–166.
- [20] D. Greenhalgh and M. Haque, A predator-prey model with disease in the prey species only, Math. Meth. Appl. Sci., 30 (2007), 911–929.
- [21] K.P. Hadeler and H.I. Freedman, Predator-prey populations with parasitic infection, J. Math. Biol., 27 (1989), 609–631.
- [22] G.S. Jiang and C.-W. Shu, Efficient implementation of weighted ENO schemes, J. Comput. Phys., 126 (1996), 202–228.
- [23] Y. Katznelson, An Introduction to Harmonic Analysis, Third Ed., Cambridge University Press, Cambridge, UK, 2004.
- [24] C.A. Kennedy and M.H. Carpenter, Additive Runge-Kutta schemes for convection-diffusionreaction equations, Appl. Numer. Math., 44 (2003), 139–181.
- [25] Y. Kuang and E. Beretta, Global qualitative analysis of a ratio-dependent predator?prey system, J. Math. Biol., 36 (1998), 389–406.
- [26] P.H. Leslie and J.C. Gower, The properties of a stochastic model for the predator-prey type of interaction between two species, *Biometrika*, 47 (1960), 219–234.
- [27] W.-M. Liu, H.W. Hethcote, and S.A. Levin, Dynamical behavior of epidemiological models with nonlinear incidence rates, J. Math. Biol., 25 (1987), 359–380.
- [28] W.-M. Liu, S.A. Levin, and Y. Iwasa, Influence of nonlinear incidence rates upon the behavior of SIRS epidemiological models, J. Math. Biol., 23 (1986), 187–204.
- [29] X.-D. Liu, S. Osher, and T. Chan, Weighted essentially non-oscillatory schemes, J. Comput. Phys., 115 (1994), 200–212.
- [30] H. Malchow, S.V. Petrovskii, and E. Venturino, Spatial Patterns in Ecology and Epidemiology: Theory, Models, and Simulation. Chapman & Hall/CRC, Boca Raton, FL, USA, 2008.
- [31] J.D. Murray, Mathematical Biology II: Spatial Models and Biomedical Applications. Third Edition. Springer, New York, 2003.
- [32] A. Okubo and S.A. Levin, Diffusion and Ecological Problems: Modern Perspectives. Second Edition, Springer-Verlag, New York, 2001.
- [33] L. Pareschi and G. Russo, Implicit-Explicit Runge-Kutta schemes and applications to hyperbolic systems with relaxation, J. Sci. Comput., 25 (2005), 129–155.
- [34] E. Rossi and V. Schleper, Convergence of a numerical scheme for a mixed hyperbolic-parabolic system in two space dimensions, ESAIM Math. Modelling Numer. Anal., 50 (2016), 475–497.
- [35] L. Sattenspiel, The Geographic Spread of Infectious Diseases: Models and Applications, Princeton Series in Theoretical and Computational Biology, Princeton University Press, 2009.
- [36] C.-W. Shu and S. Osher, Efficient implementation of essentially non-oscillatory shockcapturing schemes, II, J. Comput. Phys., 83 (1988), 32–78.
- [37] S.W. Smith, Digital Signal Processing: A Practical Guide for Engineers and Scientists. Demystifying technology series: by engineers, for engineers. Newnes, 2003.
- [38] S.H. Strogatz, Nonlinear Dynamics and Chaos: with Applications to Physics, Biology, Chemistry, and Engineering, Second Edition, CRC Press, Boca Raton, FL, 2015.
- [39] G. Q. Sun, Pattern formation of an epidemic model with diffusion, Nonlinear Dynamics, 69 (2012), 1097-1104.
- [40] P. van den Driessche, Deterministic compartmental models: extensions of basic models. In F. Brauer, P. van den Driessche, and J. Wu (Eds.), Mathematical Epidemiology, Springer-Verlag, Berlin, 2008, 147–157.
- [41] P. van den Driessche, Spatial structure: patch models. In F. Brauer, P. van den Driessche and J. Wu (Eds.), Mathematical Epidemiology, Springer-Verlag, Berlin, 2008, 179–189.

- [42] E. Vynnycky and R.E. White, An Introduction to Infectious Disease Modelling, Oxford University Press, 2010.
- [43] J. Wu, Spatial structure: partial differential equations models. In F. Brauer, P. van den Driessche and J. Wu (Eds.), Mathematical Epidemiology, Springer-Verlag, Berlin, 2008, 191– 203.
- [44] Y.-T. Zhang and C.-W Shu, ENO and WENO schemes. Chapter 5 in R. Abgrall and C.-W. Shu (eds.), Handbook of Numerical Methods for Hyperbolic Problems Basic and Fundamental Issues. Handbook of Numerical Analysis vol. 17, North Holland (2016), 103–122.
- [45] X. Zhong, Additive semi-implicit Runge-Kutta methods for computing high-speed nonequilibrium reactive flows, J. Comput. Phys., 128 (1996), 19–31.

E-mail address: rburger@ing-mat.udec.cl E-mail address: gchowell@gsu.edu E-mail address: egavilan@ing-mat.udec.cl E-mail address: mulet@uv.es E-mail address: lvillada@ubiobio.cl

Centro de Investigación en Ingeniería Matemática (Cl²MA)

PRE-PUBLICACIONES 2018

- 2018-06 JESSIKA CAMAÑO, RODOLFO RODRÍGUEZ, PABLO VENEGAS: Convergence of a lowest-order finite element method for the transmission eigenvalue problem
- 2018-07 JAVIER A. ALMONACID, GABRIEL N. GATICA, RICARDO OYARZÚA: A posteriori error analysis of a mixed-primal finite element method for the Boussinesq problem with temperature-dependant viscosity
- 2018-08 GABRIEL N. GATICA, BRYAN GOMEZ-VARGAS, RICARDO RUIZ-BAIER: Formulation and analysis of fully-mixed methods for stress-assisted diffusion problems
- 2018-09 JAY GOPALAKRISHNAN, MANUEL SOLANO, FELIPE VARGAS: Dispersion analysis of HDG methods
- 2018-10 FRANCO FAGNOLA, CARLOS M. MORA: Bifurcation analysis of a mean field laser equation
- 2018-11 DAVID MORA, IVÁN VELÁSQUEZ: A virtual element method for the transmission eigenvalue problem
- 2018-12 ALFREDO BERMÚDEZ, BIBIANA LÓPEZ-RODRÍGUEZ, RODOLFO RODRÍGUEZ, PILAR SALGADO: Numerical solution of a transient three-dimensional eddy current model with moving conductors
- 2018-13 RAIMUND BÜRGER, ENRIQUE D. FERNÁNDEZ NIETO, VICTOR OSORES: A dynamic multilayer shallow water model for polydisperse sedimentation
- 2018-14 ANTONIO BAEZA, RAIMUND BÜRGER, PEP MULET, DAVID ZORÍO: Weno reconstructions of unconditionally optimal high order
- 2018-15 VERONICA ANAYA, MOSTAFA BENDAHMANE, DAVID MORA, MAURICIO SEPÚL-VEDA: A virtual element method for a nonlocal FitzHugh-Nagumo model of cardiac electrophysiology
- 2018-16 TOMÁS BARRIOS, ROMMEL BUSTINZA: An a priori error analysis for discontinuous Lagrangian finite elements applied to nonconforming dual mixed formulations: Poisson and Stokes problems
- 2018-17 RAIMUND BÜRGER, GERARDO CHOWELL, ELVIS GAVILÁN, PEP MULET, LUIS M. VILLADA: Numerical solution of a spatio-temporal predator-prey model with infected prey

Para obtener copias de las Pre-Publicaciones, escribir o llamar a: DIRECTOR, CENTRO DE INVESTIGACIÓN EN INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C, CONCEPCIÓN, CHILE, TEL.: 41-2661324, o bien, visitar la página web del centro: http://www.ci2ma.udec.cl









Centro de Investigación en Ingeniería Matemática (CI²MA) **Universidad de Concepción**

Casilla 160-C, Concepción, Chile Tel.: 56-41-2661324/2661554/2661316http://www.ci2ma.udec.cl





