UNIVERSIDAD DE CONCEPCIÓN



Centro de Investigación en Ingeniería Matemática (CI^2MA)



A multiclass Lighthill-Whitham-Richards traffic model with a discontinuous velocity function

RAIMUND BÜRGER, CHRISTOPHE CHALONS, RAFAEL ORDOÑEZ, LUIS M. VILLADA

PREPRINT 2020-26

SERIE DE PRE-PUBLICACIONES

A MULTICLASS LIGHTHILL-WHITHAM-RICHARDS TRAFFIC MODEL WITH A DISCONTINUOUS VELOCITY FUNCTION

RAIMUND BÜRGER

Cl²MA and Departamento de Ingeniería Matemática, Universidad de Concepción, Casilla 160-C, Concepción, Chile

CHRISTOPHE CHALONS

Laboratoire de Mathématiques de Versailles Université Versailles Saint-Quentin-en-Yvelines, Versailles, France

RAFAEL ORDOÑEZ*

Cl²MA and Departamento de Ingeniería Matemática, Universidad de Concepción, Casilla 160-C, Concepción, Chile

LUIS MIGUEL VILLADA

CI²MA, Universidad de Concepción, Casilla 160-C, Concepción, Chile and Departamento de Matemáticas, Universidad del Bío-Bío, Concepción, Chile

(Communicated by the associate editor name)

ABSTRACT. The well-known Lighthill-Whitham-Richards (LWR) kinematic model of traffic flow models the evolution of the local density of cars by a nonlinear scalar conservation law. The transition between free and congested flow regimes can be described by a flux or velocity function that has a discontinuity at a determined density. A numerical scheme to handle the resulting LWR model with discontinuous velocity was proposed in [J.D. Towers, A splitting algorithm for LWR traffic models with flux discontinuities in the unknown, J. Comput. Phys., 421 (2020), article 109722]. A similar scheme is constructed by decomposing the discontinuous velocity function into a Lipschitz continuous function plus a Heaviside function and designing a corresponding splitting scheme. The part of the scheme related to the discontinuous flux is handled by a semi-implicit step that does, however, not involve the solution of systems of linear or nonlinear equations. It is proved that the whole scheme converges to a weak solution in the scalar case. The scheme can in a straightforward manner be extended to the multiclass LWR (MCLWR) model, which is defined by a hyperbolic system of N conservation laws for N driver classes that are distinguished by their preferential velocities. It is shown that the multiclass scheme satisfies an invariant region principle, that is, all densities are nonnegative and their sum does not exceed a maximum value. In the scalar and multiclass cases no flux regularization or Riemann solver is involved, and the CFL condition is not more restrictive than for an explicit scheme for the continuous part of the flux. Numerical tests for the scalar and multiclass cases are presented.

Date: October 29, 2020.

²⁰¹⁰ Mathematics Subject Classification. Primary: 65M06; Secondary: 35L45, 35L65, 76A99. Key words and phrases. Lighthill-Whitham-Richards traffic model, multiclass traffic mnodel, system of conservation laws, discontinuous flux, invariant region principle.

^{*} Corresponding author: R. Ordoñez.

1. Introduction.

1.1. Scope. The multiclass Lighthill-Whitham-Richards (MCLWR) model is a generalization of the well-known Lighthill-Whitham-Richards (LWR) model [24, 27] to multiple classes of drivers and was formulated independently by Wong and Wong [30] and Benzoni-Gavage and Colombo [1]. The model is given by the following system of conservation laws in one space dimension, where the sought unknowns are the densities $\phi_i = \phi_i(x, t)$ of vehicles of class $i, i = 1, \ldots, N$, as a function of distance x and time t [1,30]:

$$\partial_t \phi_i + \partial_x (\phi_i v_i(\phi)) = 0, \quad i = 1, \dots, N.$$
(1.1)

Here $\phi = \phi_1 + \cdots + \phi_N$ denotes the total density of vehicles. The velocity function v_i is assumed to depend on ϕ , where we assume that

$$v_i(\phi) = v_i^{\max} V(\phi), \quad i = 1, \dots, N,$$
(1.2)

where $v_1^{\max} < v_2^{\max} < \ldots < v_N^{\max}$ are the maximum velocities of the N classes of vehicles and V is a hindrance function that models the drivers' attitude to reduce speed in the presence of other cars. This function is usually assumed to be continuous and piecewise smooth on an interval $[0, \phi_{\max}]$, where $\phi_{\max} > 0$ denotes a maximum vehicle density, with

$$V(0) = 1, \quad V'(\phi) < 0, \quad V(\phi_{\max}) = 0.$$

The simplest function having all these properties is the linear interpolation $V(\phi) = 1-\phi/\phi_{\text{max}}$. However, equation (1.1) is studied herein under the assumption that V is piecewise continuous with one decreasing jump at a density value $\phi^* \in (0, \phi_{\text{max}})$, that is

$$V(\phi) = \begin{cases} V_{\rm f}(\phi) & \text{for } 0 \leqslant \phi \leqslant \phi^*, \\ V_{\rm c}(\phi) & \text{for } \phi^* < \phi \leqslant \phi_{\rm max}, \end{cases} V_{\rm f} \in C^1[0, \phi^*], \quad V_{\rm c} \in C^1[\phi^*, \phi_{\rm max}], \\ V_{\rm f}(0) = 1, \quad V_{\rm f}'(\phi) \leqslant 0 \text{ on } [0, \phi^*], \quad V_{\rm c}'(\phi) \leqslant 0 \text{ on } [\phi^*, \phi_{\rm max}], \quad V_{\rm f}(\phi_{\rm max}) = 0, \end{cases}$$
(1.3)
$$\alpha_V := V_{\rm f}(\phi^*) - V_{\rm c}(\phi^*) > 0.$$

We consider (1.1) on the domain $\Pi_T := (-L, L) \times (0, T)$, where L > 0 and T > 0, along with the initial and boundary conditions

$$\phi_{i}(x,0) = \phi_{i,0}(x) \in [0, \phi_{\max}], \quad x \in (-L,L),$$

$$\phi_{i}(-L,t) = r_{i}(t) \in [0, \phi_{\max}], \quad t \in (0,T),$$

$$\phi_{i}(L,t) = s_{i}(t) \in [0, \phi_{\max}], \quad t \in (0,T), \quad i = 1, \dots, N;$$

(1.4a)

$$\boldsymbol{\mathcal{F}}(t) \in (\boldsymbol{v}^{\max})^{\mathrm{T}} \boldsymbol{s}(t) \tilde{\boldsymbol{V}}(\boldsymbol{s}(t)), \quad t \in (0,T); \quad \boldsymbol{v}^{\max} := (v_1^{\max}, \dots, v_N^{\max})^{\mathrm{T}}.$$
(1.4b)

The non-standard boundary condition (1.4b) on the total density is required in case that $s(t) = \phi^*$, where $\mathbf{s}(t) := (s_1(t), \ldots, s_N(t))^T$ and $s(t) = s_1(t) + \cdots + s_N(t)$. This implies that we assign values to $\mathcal{F}(t)$ according to

$$\boldsymbol{\mathcal{F}}(t) = \begin{cases} (\boldsymbol{v}^{\max})^{\mathrm{T}} \boldsymbol{s}(t) V(\phi^*-) & \text{if the traffic ahead of } x = L \text{ is free-flowing,} \\ (\boldsymbol{v}^{\max})^{\mathrm{T}} \boldsymbol{s}(t) V(\phi^*+) & \text{if the traffic ahead of } x = L \text{ is congested.} \end{cases}$$
(1.5)

This assumption is motivated in a wider sense by models of phase transitions between free and congested traffic flow regimes [13, 14], and more specifically by treatments of the single-class scalar version of (1.1)-(1.4). In the scalar case the

model can be formulated following initial-boundary value problem for a scalar conservation law defined on Π_T :

$$\partial_t \phi + \partial_x f(\phi) = 0, \quad (x,t) \in \Pi_T \quad f(\phi) = v^{\max} \phi V(\phi), \phi(x,0) = \phi_0(x) \in [0,\phi_{\max}], \quad x \in (-L,L), \phi(-L,t) = r(t) \in [0,\phi_{\max}], \quad t \in (0,T), \phi(L,t) = s(t) \in [0,\phi_{\max}], \quad \mathcal{F}(t) \in \tilde{f}(s(t)) \quad t \in (0,T).$$
(1.6)

with a jump in $V(\phi)$ or equivalently, in the flux $f(\phi)$, see [28,29], where \tilde{f} denotes the multivalued version of f and $\mathcal{F}(t) \in \tilde{f}(s(t))$ represents the non-standard boundary condition of the flux discontinuity, see [28].

It is the purpose of the present contribution to introduce a numerical scheme for the MCLWR model with discontinuous flux (1.1)-(1.3) that is based on the available treatment [28] of the scalar model (1.6). The scalar version of the scheme slightly differs from that of [28] but we prove that it produces approximations that also converge to a weak solution. Numerical experiments provide evidence that it approximates the same solutions as the scheme of [28]. In the multiclass case we prove satisfaction of an invariant region principle, that is, numerical solutions assume values in

$$\mathcal{D} := \left\{ (\phi_1, \dots, \phi_N)^{\mathrm{T}} \in \mathbb{R}^N : \phi_1 \ge 0, \dots, \phi_N \ge 0, \, \phi = \phi_1 + \dots + \phi_N \leqslant \phi_{\max} \right\}$$

under corresponding assumptions on the initial and boundary data.

1.2. **Related work.** The MCLWR model (1.1) has been studied intensively in recent years. The system (1.1), (1.2) has some interesting properties and in particular admits a separable entropy function for an arbitrary number of driver classes. We refer to [1, 2, 5, 7, 9-11, 19, 20, 30-36] for numerical and analytical treatments and emphasize that to our knowledge a velocity function discontinuous in the unknowns has not been considered so far for the MCLWR model.

Conservation laws with discontinuous flux function arise in many physical applications including flow in porous media [22], sedimentation [8, 18], and the LWR traffic model [25, 29]. Here we limit the discussion to analyses where the flux is a discontinuous function of the unknown (as opposed to the more widely studied discontinuous dependence on spatial position). This property implies that standard numerical methods cannot be applied directly due to the presence of waves that travel at infinite speed, namely so-called zero waves. These waves carry information about the flux but this value is transported instantaneously, which excludes applying explicit schemes due to the lack of regularity of the flux. Gimse [21] was the first to present a solution to this problem. He studied a conservation law where the flux function has a single jump. He discussed the existence of the zero wave, generalized the method of convex hull construction, and solved the Riemann problem using a front tracking algorithm.

Carrillo [12] studied conservation laws with a discontinuous flux function where the flux is allowed to have discontinuities on a finite subset of real numbers. The proof of existence of solutions is based on the comparison principle and an entropy inequality involving a version of semi-Kružkov entropies. Dias and Figueira [15] studied a related problem by using a mollification technique to smooth out discontinuities. They showed that solutions to a suitably regularized problem converge to solutions of the original problem in the limit. They also defined the notions of weak solution and weak entropy solution. The mollification technique was implemented in [16, 17]. Moreover, Dias and Figueira [16] proposed a numerical scheme for Riemann problem. Specifically, they introduced a procedure to obtain a new Lipschitz continuous flux function with the same lower convex envelope of the original flux, and then a standard Lax-Friedrichs method is employed.

Martin and Vovelle [26] considered the problem of numerical approximation in the Cauchy-Dirichlet problem for a scalar conservation law with a flux function having finitely many discontinuities. The well-posedness of this problem had been proved by Carrillo. An implicit finite volume scheme is constructed in [26] and Newton's method is employed to solve the resulting system of nonlinear equations. Furthermore, convergence to the unique entropy solution is shown.

Lu et al. [25] explicitly constructed the entropy solutions for the LWR traffic flow model with a piecewise quadratic flow-density relationship. Their approach is based on constructions of entropy solutions to a sequence of approximate problems in which the flow-density relationship is continuous but tends to the discontinuous flux when a small parameter in this sequence tends to zero.

Bulíček et al. [3] introduced new concepts of entropy weak and measure-valued solutions that are consistent with the standard ones if the flux is continuous. They identified a given discontinuous flux function with a continuous curve that consists of the graph of this flux and abscissae that fill the jumps. Consequently, instead of a discontinuous flux function of the unknown, they deal with an implicit relation that represents a curve. One has one degree of freedom to set up the "optimal" unknown (independent variable). These ideas are combined in [4], where the authors treat the case of a flux function discontinuous in spatial position and the unknown. Through appropriate estimates for entropy measure-valued solutions well-posedness is shown.

Wiens et al. [29] applied Dias and Figueira's mollification approach to solving a conservation law with a piecewise linear flux function in which there is a single discontinuity at a critical point. They introduced a mollified function and then the analytical solution to the corresponding Riemann problem is derived in the limit. Furthermore they constructed a Riemann solver that forms the basis for a highresolution finite volume scheme of Godunov type and used an alternate approach that eliminates the severe CFL constraint by incorporating the effect of zero waves directly into the local Riemann solver.

Towers [28] presented a finite difference scheme that implement a splitting consistent with the decomposition the flux f(u) = p(u) + g(u), where p is a Lipschitz continuous function and g is a function of Heaviside type that includes the jumps of f. The scheme has the form (see [28, Eq. (3.11)])

$$\begin{cases} \begin{cases} U_j^{n+1/2} = \tilde{G}^{-1} (U_j^n - \lambda g_{j+1}^{n+1/2}), & j = M, M-1, \dots, 1, \\ g_j^{n+1/2} = (U_j^{n+1/2} - U_j^n + \lambda g_{j+1}^{n+1/2})/\lambda, & j = M, M-1, \dots, 1, \\ U_j^{n+1} = U_j^{n+1/2} - \lambda \Delta_{-} \tilde{p} (U_{j+1}^{n+1/2} - U_j^{n+1/2}), & j = 1, \dots, M, \end{cases}$$
(1.7)

which can be written in conservation form as follows:

$$U_{j}^{n+1/2} = U_{j}^{n} - \lambda \left(g_{j+1}^{n+1/2} - g_{j}^{n+1/2} \right),$$

$$U^{n+1} = U_{j}^{n+1/2} - \lambda \Delta_{-} \tilde{p} \left(U_{j+1}^{n+1/2}, U_{j}^{n+1/2} \right).$$

The first part of the scheme is implicit and consistent with $u_t + g(u)_x = 0$, but the resulting equations can be solved by evaluation of a piecewise linear function. Hence, an iterative solver like Newton's method is not required. The second part of the scheme is consistent with $u_t + p(u)_x = 0$ and is explicit, and can be solved



FIGURE 1. (a) Piecewise continuous velocity function $V(\phi)$ with discontinuity at $\phi = \phi^*$, (b) continuous and discontinuous portions $p_V(\phi)$ (solid line) and $g_V(\phi)$ (dashed line).

by any scheme suitable for a scalar conservation law with Lipschitz continuous flux. Towers [28] focused on the Godunov flux for the explicit part but also presented a simple flux-limited Lax-Wendroff-type modification to the Godunov scheme.

1.3. Outline of the paper. The remainder of the paper is organized as follows. In Section 2 we present a numerical scheme for the LWR traffic flow model. We first introduce some assumptions and the notion of weak solution in Section 2.1. Next, Section 2.2 is devoted to the presentation of our scheme for the scalar case (N = 1) and we imposed the appropriate CFL condition. Then, in Section 2.3, we prove that under the CFL condition it satisfies uniform L^{∞} and TVD properties. Moreover we prove some kind time continuity estimates and to the end this section we prove the convergence of our numerical solution converge to weak solution in sense of Definition 2.1. In Section 3 we extend the algorithm to the multiclass case (N > 1) and prove that the scheme preserves the invariant region \mathcal{D} . In Section 4 we present several numerical examples to confirm all the results mentioned before. Section 5 collects some conclusions.

2. Construction of the numerical scheme in the scalar case. Before describing the numerical scheme we introduce some assumptions and the definition of weak solutions proposed in [16], which is employed herein.

2.1. **Preliminaries.** To outline the basic idea, and to make the comparison with [28] transparent, we define the functions

$$g_V(\phi) := \alpha_V H(\phi^* - \phi), \quad p_V(\phi) := V(\phi) - g_V(\phi),$$
 (2.1)

where p_V is a Lipschitz continuous, piecewise smooth and decreasing function, while $g_V(\phi)$ is a non-negative and decreasing function, see Figure 1. Furthermore, as in [28], we can equivalently specify

$$\mathcal{G}(t) \in \tilde{g}_V(s(t)), \tag{2.2}$$

where \tilde{g}_V denotes the multivalued version of g_V . With respect to the initial and boundary data we assume that the initial density function ϕ_0 satisfies

 $\phi_0(x) \in [0, \phi_{\max}] \quad \text{for } x \in (-L, L), \quad \phi_0 \in BV([-L, L]), \quad g_V(\phi_0) \in BV([-L, L]).$

The boundary functions r and s are assumed to satisfy

$$r(t), s(t) \in [0, \phi_{\max}] \text{ for } t \in [0, T], r, s \in BV([0, T]).$$

We also assume that $\mathcal{G}(t) \in [0, \alpha_V]$ for all $t \in [0, T]$, and $\mathcal{G} \in BV([0, T])$.

Definition 2.1 (Weak solution [16]). A function $\phi \in L^{\infty}(\Pi_T)$ is said to be a *weak solution* to the initial-boundary value problem (1.6) if there exists a function $q \in L^{\infty}(\Pi_T)$ satisfying $q(x,t) \in \tilde{f}(\phi(x,t))$ a.e. such that for all test functions $\psi \in C_0^1([-L, L] \times [0, T)),$

$$\int_0^T \int_{-L}^L (\phi \psi_t + q \psi_x) \, \mathrm{d}x \, \mathrm{d}t + \int_{-L}^L \phi_0(x) \psi(x,0) \, \mathrm{d}x = 0.$$

2.2. Numerical scheme. The domain Π_T is discretized as follows. We choose a partition $\{I_j\}_{j=1}^M$ of [-L, L] composed of uniform cells $I_j = [x_{j-1/2}, x_{j+1/2}]$, where $x_{j+1/2} = x_j + \Delta x/2$, that are centered in x_j and have length $|I_j| = \Delta x = 2L/M$. Then, for $\Delta t > 0$, we let $t^n = n\Delta t$ for $n = 0, \ldots, \mathcal{N}$, where \mathcal{N} is an integer such that $T \in [t^{\mathcal{N}}, t^{\mathcal{N}} + \Delta t)$. The unknowns ϕ_j^n approximate the cell average of the exact solution $\phi(\cdot, t^n)$ in the cell I_j . The initial condition is discretized by

$$\phi_j^0 = \frac{1}{\Delta x} \int_{I_j} \phi_0(x) \,\mathrm{d}x, \quad j = 1, \dots, M,$$

and the boundary conditions with $\mathcal{F}(t) \in \tilde{f}(s)$ are discretized as follows:

$$\begin{split} \phi_{0}^{n+1/2} &= \phi_{0}^{n} = r(t^{n}) = t^{n}, \quad \phi_{M+1}^{n+1/2} = \phi_{M+1}^{n} = s(t^{n}) = s^{n}, \\ r^{n} &\in [0, \phi_{\max}], \quad s^{n} \in [0, \phi_{\max}], \quad g_{M+1}^{n+1/2} \in [0, \alpha_{V}], \\ g_{M+1}^{n+1/2} &= g_{M+1}^{n} = \mathcal{G}(s^{n}) \\ &= \begin{cases} \alpha_{V} & \text{if } s^{n} < \phi^{*}, \\ \alpha_{V} & \text{if } s^{n} = \phi^{*} \text{ and traffic ahead of } x = L \text{ is free-flowing,} \\ 0 & \text{if } s^{n} = \phi^{*}. \end{cases} \end{split}$$
(2.3)

Before proposing our scheme we recall that the basic idea of a splitting scheme consists in solving within each time step, first the PDE

$$\partial_t \phi + \partial_x \left(v^{\max} \phi g_V(\phi) \right) = 0, \tag{2.4}$$

followed by the solution of the conservation law with continuous flux

$$\partial_t \phi + \partial_x \left(v^{\max} \phi p_V(\phi) \right) = 0. \tag{2.5}$$

Note that in the scalar case the constant v^{\max} is immaterial. For the remainder of the analysis of the scalar case we assume that t or x are rescaled so that $v^{\max} = 1$.

Based on the form of the flux function of equations (2.4) and (2.5) and the properties of the functions g_V and p_V , we may write a numerical scheme for (1.6)

MULTICLASS TRAFFIC WITH DISCONTINUOUS VELOCITY



FIGURE 2. (a) function $z \mapsto \tilde{G}_V(z; \phi)$ given by (2.9a) with $\lambda v^{\max} = 1/2$, $\alpha_V = 0.3$, and $\phi = 0.8$, (b) its inverse $z \mapsto \tilde{G}_V^{-1}(z; \phi)$ given by (2.9b).

that is motivated by Scheme 4 of [6] in the following form:

$$\phi_{j}^{n+1/2} = \phi_{j}^{n} - \lambda \left(\phi_{j}^{n} g_{V,j+1}^{n+1/2} - \phi_{j-1}^{n} g_{V,j}^{n+1/2} \right),$$

$$\phi_{j}^{n+1} = \phi_{j}^{n+1/2} - \lambda \left(\phi_{j}^{n+1/2} p_{V} \left(\phi_{j+1}^{n+1/2} \right) - \phi_{j-1}^{n+1/2} p_{V} \left(\phi_{j}^{n+1/2} \right) \right), \qquad (2.6)$$

$$j = 1, \dots, M.$$

The first half-step in (2.6) is semi-implicit and is consistent with (2.4) whereas the second half-step is explicit and consistent with (2.5).

In order to evaluate the first line in (2.6), we start by computing the values $g_{V,j}^{n+1/2}$ from j = M + 1 to j = 1 (in decreasing order). This is motivated by the following argument, where we start from the semi-implicit equation

$$\phi_j^{n+1/2} = \phi_j^n - \lambda \left(\phi_j^n g_V \left(\phi_{j+1}^{n+1/2} \right) - \phi_{j-1}^n g_V \left(\phi_j^{n+1/2} \right) \right)$$
(2.7)

along with a known value $\mathcal{G}(\phi_{M+1}^{n+1/2})$ arising from the boundary condition. Next, we write $g_V(\phi_{j+1}^{n+1/2})$ as $g_{V,j+1}^{n+1/2}$ and then rearrange (2.7) as

$$\phi_j^{n+1/2} - \lambda \phi_{j-1}^n g_V (\phi_j^{n+1/2}) = \phi_j^n - \lambda \phi_j^n g_{V,j+1}^{n+1/2}.$$
(2.8)

Let us now define the function

$$G_V(z;\phi) := z - \lambda \phi g_V(z), \quad z, \phi \in [0, \phi_{\max}]$$

along with its multivalued version (with respect to z) $\tilde{G}_V(\cdot;\phi)$. Then \tilde{G}_V is strictly increasing and has a unique inverse $z \mapsto \tilde{G}_V^{-1}(z;\phi)$, see Figure 2. Explicitly, we get

$$\tilde{G}_{V}(z;\phi) := \begin{cases} z - \lambda \alpha_{V}\phi & \text{for } z \in [0, \phi^{*}), \\ [\phi^{*} - \lambda \alpha_{V}\phi, \phi^{*}] & \text{for } \phi = \phi^{*}, \\ z & \text{for } z \in [\phi^{*}, \phi_{\max}], \end{cases}$$

$$\tilde{G}_{V}^{-1}(z;\phi) := \begin{cases} z + \lambda \alpha_{V}\phi & \text{for } z \in [-\lambda \alpha_{V}\phi, \phi^{*} - \lambda \alpha_{V}\phi), \\ \phi^{*} & \text{for } z \in [\phi^{*} - \lambda \alpha_{V}\phi, \phi^{*}], \\ z & \text{for } z \in [\phi^{*}, \phi_{\max}]. \end{cases}$$
(2.9a)
$$\tilde{G}_{V}^{-1}(z;\phi) := \begin{cases} z + \lambda \alpha_{V}\phi & \text{for } z \in [-\lambda \alpha_{V}\phi, \phi^{*}], \\ \phi^{*} & \text{for } z \in [\phi^{*}, \phi_{\max}]. \end{cases}$$

Consequently, we may express (2.8) as

$$\tilde{G}_V(\phi_j^{n+1/2};\phi_{j-1}^n) = \phi_j^n - \lambda \phi_j^n g_{V,j+1}^{n+1/2},$$

which allows us to obtain $\phi_j^{n+1/2}$ by applying $\tilde{G}_V^{-1}(z;\phi)$ to both sides, that is

$$\phi_j^{n+1/2} = \tilde{G}_V^{-1} \left(\phi_j^n - \lambda \phi_j^n g_{V,j+1}^{n+1/2}; \phi_{j-1}^n \right).$$
(2.10)

Now that $\phi_j^{n+1/2}$ is available, we solve for $g_{V,j}^{n+1/2}$ the equation

$$\phi_j^{n+1/2} = \phi_j^n - \lambda \left(\phi_j^n g_{V,j+1}^{n+1/2} - \phi_{j-1}^n g_{V,j}^{n+1/2} \right), \tag{2.11}$$

provided that $\phi_{j-1}^n > 0$. If $\phi_{j-1}^n = 0$, we define directly

$$g_{V,j}^{n+1/2} = g_V(\phi_j^{n+1/2})$$

The numerical scheme can be summarized in Algorithm 2.1:

Algorithm 2.1 (BCOV scheme, scalar case).

$$\begin{array}{l} \mbox{Input: approximate solution vector } \{\phi_{j}^{n}\}_{j=1}^{M} \mbox{ for } t = t^{n} \\ g_{V,M+1}^{n+1/2} \leftarrow \mathcal{G}(\phi_{M+1}^{n+1/2}) \mbox{ (using (2.3))} \\ \mbox{do } j = M, M-1, \dots, 1 \\ \phi_{j}^{n+1/2} \leftarrow \tilde{G}_{V}^{-1}(\phi_{j}^{n} - \lambda \phi_{j}^{n}g_{V,j+1}^{n+1/2}; \phi_{j-1}^{n}) \\ \mbox{if } \phi_{j-1}^{n} \neq 0 \mbox{ then} \\ g_{V,j}^{n+1/2} \leftarrow \frac{\phi_{j}^{n+1/2} - \phi_{j}^{n} + \lambda g_{V,j+1}^{n+1/2} \phi_{j}^{n}}{\lambda \phi_{j-1}^{n}} \\ \mbox{else} \\ g_{V,j}^{n+1/2} \leftarrow g_{V}(\phi_{j}^{n+1/2}) \\ \mbox{endif} \\ \mbox{enddo} \\ \mbox{do } j = 1, 2, \dots, M \\ \phi_{j}^{n+1} \leftarrow \phi_{j}^{n+1/2} - \lambda \big(\phi_{j}^{n+1/2} p_{V}(\phi_{j+1}^{n+1/2}) - \phi_{j-1}^{n+1/2} p_{V}(\phi_{j}^{n+1/2}) \big) \\ \mbox{enddo} \\ \mbox{Output: approximate solution vector } \{\phi_{j}^{n+1}\}_{j=1}^{M} \mbox{ for } t = t^{n+1} = t^{n} \end{array}$$

Next, we demonstrate that the numerical scheme (2.11) is consistent with (2.4).

 $+\Delta t$

Lemma 2.1. Assume that $\phi_{V,j}^{n+1/2} \in [0, \phi_{\max}]$ for all j. Then $g_{V,j}^{n+1/2} \in \tilde{g}_V(\phi_j^{n+1/2})$ for all j. In particular $g_{V,j}^{n+1/2} \in [0, \alpha_V]$ for all j.

Proof. Let us first assume that $\phi_{j-1} = 0$. Then the result follows from the definition of the function $g_V(z)$ and the corresponding assignment to $g_{V,j}^{n+1/2}$ in Algorithm 2.1. If $\phi_{j-1} \neq 0$, then (2.10) and (2.9) imply that

$$\phi_j^{n+1/2} - \lambda \phi_{j-1}^n g_{V,j}^{n+1/2} \in \tilde{G}_V(\phi_j^{n+1/2}; \phi_{j-1}^n).$$

Therefore, by a straightforward case-by-case study (of the cases arising in (2.9)) we conclude that $g_{V,j}^{n+1/2} \in \tilde{g}_V(\phi_j^{n+1/2})$.

Now, to derive CFL conditions, we write the scheme (2.6) in incremental form

$$\phi_j^{n+1/2} = \phi_j^n + C_{g,j+1/2}^{n+1/2} \Delta_+ \phi_j^{n+1/2} - D_{g,j-1/2}^{n+1/2} \Delta_- \phi_j^n, \qquad (2.12a)$$

$$\phi_j^{n+1} = \phi_j^{n+1/2} + C_{p,j+1/2}^{n+1/2} \Delta_+ \phi_j^{n+1/2} - D_{p,j-1/2}^{n+1/2} \Delta_- \phi_j^{n+1/2}$$
(2.12b)

with the spatial difference operators $\Delta_+ V_j := V_{j+1} - V_j$ and $\Delta_- V_j := V_j - V_{j-1}$ and the incremental coefficients

$$\begin{split} C_{g,j+1/2}^{n+1/2} &:= \begin{cases} \lambda \phi_j^n \frac{g_V(\phi_j^{n+1/2}) - g_V(\phi_{j+1}^{n+1/2})}{\phi_{j+1}^{n+1/2} - \phi_j^{n+1/2}} & \text{if } \phi_{j+1}^{n+1/2} \neq \phi_j^{n+1/2}, \\ 0 & \text{otherwise}, \end{cases} \\ D_{g,j-1/2}^{n+1/2} &:= \lambda g_V(\phi_j^{n+1/2}), \\ C_{p,j+1/2}^{n+1/2} &:= \begin{cases} \lambda \phi_j^{n+1/2} \frac{p_V(\phi_j^{n+1/2}) - p_V(\phi_{j+1}^{n+1/2})}{\phi_{j+1}^{n+1/2} - \phi_j^{n+1/2}} & \text{if } \phi_{j+1}^{n+1/2} \neq \phi_j^{n+1/2}, \\ 0 & \text{otherwise}, \end{cases} \\ D_{p,j-1/2}^{n+1/2} &:= \lambda p_V(\phi_j^{n+1/2}). \end{split}$$

To have an L^{∞} estimate (Lemma 2.2 below) and the Total Variation Decreasing (TVD) property (Lemma 2.3 below) sufficient conditions are

$$0 \leqslant D_{p,j-1/2}^{n+1/2}, C_{p,j+1/2}^{n+1/2} \leqslant \frac{1}{2}, \quad C_{g,j+1/2}^{n+1/2} \geqslant 0, \quad 0 \leqslant D_{g,j-1/2}^{n+1/2} \leqslant 1 \quad \text{for all } j$$

First, we observe the following fact about \tilde{g}_V . If $z_1, z_2 \in [0, \phi_{\max}]$ and $z_1 \neq z_2$, then

$$g_{V,1} \in \tilde{g}_V(z_1), \ g_{V,2} \in \tilde{g}_V(z_2) \Longrightarrow \frac{g_{V,2} - g_{V,1}}{z_2 - z_1} \leqslant 0.$$
 (2.13)

This property and Lemma 2.1 imply that

$$D_{g,j-1/2}^{n+1/2}, C_{g,j+1/2}^{n+1/2} \ge 0$$
 for all j .

Next, the properties of the function p_V ensure that

$$C_{p,j+1/2}^{n+1/2}, D_{p,j-1/2}^{n+1/2} \ge 0$$
 for all j

Finally, to enforce the inequalities

$$D_{p,j-1/2}^{n+1/2}, C_{p,j+1/2} \leqslant \frac{1}{2}$$
 and $D_{g,j-1/2}^{n+1/2} \leqslant 1$ for all j ,

we impose the CFL conditions

$$\lambda \Big(\phi_{\max} \max_{1 \leq j \leq M} \left| p'_V(\phi_j) \right| + \max_{1 \leq j \leq M} p_V(\phi_j) \Big) \leq \frac{1}{2}, \qquad \lambda \alpha_V \leq 1.$$
(2.14)

2.3. Convergence of the scalar scheme. The goal is to prove convergence of approximate solution to a weak solution of (1.6). The discrete solutions $\{\phi_j^{n+1/2}\}$ constructed via the scheme (2.6) are extended to the whole domain Π_T by defining the piecewise constant function

$$\phi^{\Delta}(x,t) = \sum_{n=0}^{N} \sum_{j=1}^{M} \chi_j(x) \chi^n(t) \phi_j^{n+1/2}$$
(2.15)

where $\Delta = (\Delta x, \Delta t)$ and $\chi_j(x)$ and $\chi^n(t)$ are the characteristic functions of cell I_j and the time interval $[t^n, t^n + \Delta t)$, respectively. The ratio $\lambda = \Delta t / \Delta x$ is always kept constant, so the limits $\Delta t \to 0$, $\Delta x \to 0$, and $\Delta \to \mathbf{0}$ are equivalent.

We start by proving an L^{∞} estimate on ϕ^{Δ} . In the remainder of this section it is always assumed that the CFL condition (2.14) is in effect.

Lemma 2.2. If $\phi_j^0 \in [0, \phi_{\max}]$ for $j = 1, \ldots, M$, then

 $\phi_j^n, \phi_j^{n+1/2} \in [0, \phi_{\max}] \text{ for all } j = 1, \dots, M \text{ and } n = 1, \dots, \mathcal{N}.$ (2.16)

Proof. Taking n = 0 and j = M in (2.10) yields

$$\phi_M^{1/2} = \tilde{G}_V^{-1} \left(\phi_M^0 - \lambda \phi_M^0 g_{V,M+1}^{1/2}; \phi_{M-1}^0 \right).$$
(2.17)

The boundary condition $g_{V,M+1}^{1/2} = \mathcal{G}(t^0) \subseteq [0, \alpha_V]$ together with the assumption implies that

$$-\lambda \alpha_V \phi_M^0 \leqslant \phi_M^0 - \lambda \phi_M^0 g_{V,M+1}^{1/2} \leqslant \phi_{\max}.$$

Since $\tilde{G}_V^{-1}(\cdot;\phi)$ is a nondecreasing function and maps $[-\lambda\alpha_V\phi,\phi_{\max}]$ onto $[0,\phi_{\max}]$, (2.17) implies that $\phi_M^{1/2} \in [0,\phi_{\max}]$. It follows from (2.1) that $g_{V,M}^{1/2} \in [0,\alpha_V]$. Reasoning in this way for $j = M-1, M-2, \ldots, 1$ yields $\phi_j^{1/2} \in [0,\phi_{\max}]$ for $j = 1,\ldots,M$. Since $\phi_0^{1/2}, \phi_{M+1}^{1/2} \in [0,\phi_{\max}]$ by (2.3), and taking into account (2.12), we find that ϕ_j^1 is a convex combination of $\phi_{j-1}^{1/2}, \phi_j^{1/2}$ and $\phi_{j+1}^{1/2}$. Thus, $\phi_j^1 \in [0,\phi_{\max}]$ for $j = 1,\ldots,M$. Repeating this argument inductively for $n = 1,\ldots,N$ we obtain (2.16).

Lemma 2.3. The discrete approximate solutions generated by the scheme (2.12) satisfy the following spatial variation bounds:

$$\sum_{j=0}^{M} \left| \phi_{j+1}^{n} - \phi_{j}^{n} \right| \leq \mathrm{TV}(\phi_{0}) + \mathrm{TV}(r) + \mathrm{TV}(s),$$

$$\sum_{j=0}^{M} \left| \phi_{j+1}^{n+1/2} - \phi_{j}^{n+1/2} \right| \leq \mathrm{TV}(\phi_{0}) + \mathrm{TV}(r) + \mathrm{TV}(s).$$
(2.18)

Proof. Applying the operator Δ_+ to (2.12a) and rearranging yields

$$(1 + C_{g,j+1/2}^{n+1/2}) \Delta_+ \phi_j^{n+1/2} = (1 - D_{g,j+1/2}^{n+1/2}) \Delta_+ \phi_j^n + C_{g,j+3/2}^{n+1/2} \Delta_+ \phi_{j+1}^{n+1/2} + D_{g,j-1/2}^{n+1/2} \Delta_+ \phi_{j-1}^n$$

Taking absolute values, summing over j = 1, ..., M - 1 and using (2.14) we get

$$\begin{split} &\sum_{j=1}^{M-1} \left(1 + C_{g,j+1/2}^{n+1/2} \right) \left| \Delta_{+} \phi_{j}^{n+1/2} \right| \\ &\leqslant \sum_{j=1}^{M-1} \left(1 - D_{g,j+1/2}^{n+1/2} \right) \left| \Delta_{+} \phi_{j}^{n} \right| + \sum_{j=1}^{M-1} C_{g,j+3/2}^{n+1/2} \left| \Delta_{+} \phi_{j+1}^{n+1/2} \right| \\ &+ \sum_{j=1}^{M-1} D_{g,j-1/2}^{n+1/2} \left| \Delta_{+} \phi_{j-1}^{n} \right|. \end{split}$$

Cancelling telescoping terms we obtain

$$\sum_{j=1}^{M-1} \left| \Delta_{+} \phi_{j}^{n+1/2} \right| + C_{g,3/2}^{n+1/2} \left| \Delta_{+} \phi_{1}^{n+1/2} \right|$$

$$\leq \sum_{j=1}^{M-1} \left| \Delta_{+} \phi_{j}^{n} \right| - D_{g,M-1/2}^{n+1/2} \left| \Delta_{+} \phi_{M-1}^{n} \right| + C_{g,M+1/2}^{n+1/2} \left| \Delta_{+} \phi_{M}^{n+1/2} \right|$$

$$+ D_{g,1/2}^{n+1/2} \left| \Delta_{+} \phi_{0}^{n+1/2} \right|.$$
(2.19)

The boundary condition implies

$$\Delta_{+}\phi_{0}^{n+1/2} = \left(1 - D_{g,1/2}^{n+1/2}\right)\Delta_{+}\phi_{0}^{n} + C_{g,3/2}^{n+1/2}\Delta_{+}\phi_{1}^{n+1/2},$$
$$\left(1 + C_{g,M+1/2}^{n+1/2}\right)\Delta_{+}\phi_{M}^{n+1/2} = \Delta_{+}\phi_{M}^{n} + D_{g,M-1/2}^{n+1/2}\Delta_{+}\phi_{1}^{n+1/2}.$$

After taking absolute values in the two previous equations, we get

$$\left|\Delta_{+}\phi_{0}^{n+1/2}\right| \leq \left(1 - D_{g,1/2}^{n+1/2}\right) \left|\Delta_{+}\phi_{0}^{n}\right| + C_{g,3/2}^{n+1/2} \left|\Delta_{+}\phi_{1}^{n+1/2}\right|,$$

$$\left(1 + C_{g,M+1/2}^{n+1/2}\right) \left|\Delta_{+}\phi_{M}^{n+1/2}\right| \leq \left|\Delta_{+}\phi_{M}^{n}\right| + D_{g,M-1/2}^{n+1/2} \left|\Delta_{+}\phi_{M-1}^{n+1/2}\right|.$$
(2.20)

From (2.19) and (2.20)

$$\sum_{j=0}^{M} \left| \Delta_{+} \phi_{j}^{n+1/2} \right| \leqslant \sum_{j=0}^{M} \left| \Delta_{+} \phi_{j}^{n} \right|.$$
(2.21)

Reasoning in the same way as the proof of Lemma 5.2 in [28] we find that

$$\sum_{j=0}^{M} \left| \Delta_{+} \phi_{j}^{n+1} \right| \leq \sum_{j=0}^{M} \left| \Delta_{+} \phi_{j}^{n} \right| + \left| r^{n+1} - r^{n} \right| + \left| s^{n+1} - s^{n} \right|.$$

It follows by induction that

$$\sum_{j=0}^{M} \left| \phi_{j+1}^{n} - \phi_{j}^{n} \right| \leq \mathrm{TV}(\phi_{0}) + \mathrm{TV}(r) + \mathrm{TV}(s).$$
(2.22)

From (2.21) and (2.22) we get (2.18).

Now, we prove some time continuity estimates. The proof of the first of them is very similar to that of [28, Lemma 5.5], so we omit the details.

Lemma 2.4. The following discrete L^1 time continuity estimate holds for $n \ge 0$:

$$\sum_{j=0}^{M+1} |\phi_j^{n+1} - \phi_j^{n+1/2}| \leq \Omega_1, \quad \Omega_1 := \mathrm{TV}(\phi_0) + \mathrm{TV}(r) + \mathrm{TV}(s) + 2\phi_{\max}.$$

Lemma 2.5. The following estimate holds:

$$\sum_{j=1}^{M} \left| \phi_{j}^{1/2} - \phi_{j}^{0} \right| \leqslant \Omega_{2}, \quad \Omega_{2} := \sum_{j=1}^{M} \left| g_{V} \left(\phi_{j+1}^{0} \right) - g_{V} \left(\phi_{j}^{0} \right) \right| + \operatorname{TV}(\phi_{0}) + \phi_{\max}. \quad (2.23)$$

Proof. We define $g_{V,j}^0 = g_V(\phi_j^0)$. The first equation in (2.6) with n = 0 implies $\phi^{1/2} - \phi^0$

$$\begin{split} \varphi_{j} &= \varphi_{j} \\ &= \lambda \phi_{j-1}^{0} \left(g_{V,j}^{1/2} - g_{V,j}^{0} \right) - \lambda \phi_{j}^{0} \left(g_{V,j+1}^{1/2} - g_{V,j+1}^{0} \right) - \lambda \phi_{j}^{0} \left(\Delta_{+} g_{V,j}^{0} \right) - \lambda g_{V,j}^{0} \left(\Delta_{+} \phi_{j-1}^{0} \right). \end{split}$$
Thus

Т nus

$$\phi_j^{1/2} - \phi_j^0 - \lambda \phi_{j-1}^0 \left(g_{V,j}^{1/2} - g_{V,j}^0 \right)$$

$$= \lambda \phi_j^0 \left(g_{V,j+1}^{1/2} - g_{V,j+1}^0 \right) - \lambda \left(\phi_j^0 \Delta_+ g_{V,j}^0 + g_{V,j}^0 \Delta_+ \phi_{j-1}^0 \right).$$

$$(2.24)$$

Taking absolute values in (2.24) and using (2.13) we find that

$$\begin{split} & \left| \phi_j^{1/2} - \phi_j^0 \right| + \lambda \phi_{j-1}^0 \left| g_{V,j}^{1/2} - g_{V,j}^0 \right| \\ & \leqslant \lambda \phi_j^0 \left| g_{V,j+1}^{1/2} - g_{V,j+1}^0 \right| + \lambda \phi_j^0 \left| \Delta_+ g_{V,j}^0 \right| + \lambda g_{V,j}^0 \left| \Delta_+ \phi_{j-1}^0 \right|. \end{split}$$

Summing over j = 1, ..., M and cancelling telescoping terms yields

$$\sum_{j=1}^{M} |\phi_{j}^{1/2} - \phi_{j}^{0}| + \lambda \phi_{0}^{0} |g_{V,1}^{1/2} - g_{V,1}^{0}|$$

$$\leq \lambda \phi_{M}^{0} |g_{V,M+1}^{1/2} - g_{V,M+1}^{0}| + \lambda \sum_{j=1}^{M} \phi_{j}^{0} |\Delta_{+} g_{V,j}^{0}| + \lambda \sum_{j=1}^{M} g_{V,j}^{0} |\Delta_{+} \phi_{j-1}^{0}|.$$

$$(2.25)$$

Applying the boundary condition in (2.25), Lemmas 2.1, 2.2, and the CFL condition (2.14) we get (2.23).

Lemma 2.6. There exists a constant Ω_3 that is independent of Δ such that the following time continuity estimate holds:

$$\sum_{j=0}^{M+1} \left| \phi_j^{n+1/2} - \phi_j^{n-1/2} \right| \leqslant \Omega_3 \quad \text{for } n \ge 1.$$

Proof. For $n \ge 2$ and subtracting from the first half-step of (2.6) the corresponding formula for $\phi_j^{n-1/2}$ and rearranging terms we get

$$\begin{split} \phi_j^{n+1/2} &- \phi_j^{n-1/2} - \lambda \phi_{j-1}^{n-1} \left(g_{V,j}^{n+1/2} - g_{V,j}^{n-1/2} \right) \\ &= \left(1 - \lambda g_{V,j+1}^{n+1/2} \right) \left(\phi_j^n - \phi_j^{n-1} \right) + \lambda g_{V,j}^{n+1/2} \left(\phi_{j-1}^n - \phi_{j-1}^{n-1} \right) - \lambda \phi_j^{n-1} \left(g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2} \right). \end{split}$$

Taking absolute values and applying the CFL condition (2.14) yields

$$\begin{split} & \left|\phi_{j}^{n+1/2} - \phi_{j}^{n-1/2} - \lambda \phi_{j-1}^{n-1} \left(g_{V,j}^{n+1/2} - g_{V,j}^{n-1/2}\right)\right| \\ & \leqslant \left(1 - \lambda g_{V,j+1}^{n+1/2}\right) \left|\phi_{j}^{n} - \phi_{j}^{n-1}\right| + \lambda g_{V,j}^{n+1/2} \left|\phi_{j-1}^{n} - \phi_{j-1}^{n-1}\right| \\ & + \left|\lambda \phi_{j}^{n-1} \left(g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2}\right)\right|. \end{split}$$

From (2.13) we get

$$\begin{aligned} \left|\phi_{j}^{n+1/2} - \phi_{j}^{n-1/2}\right| + \left|\lambda\phi_{j-1}^{n-1}\left(g_{V,j}^{n+1/2} - g_{V,j}^{n-1/2}\right)\right| \\ &\leqslant \left(1 - \lambda g_{V,j+1}^{n+1/2}\right) \left|\phi_{j}^{n} - \phi_{j}^{n-1}\right| + \lambda g_{V,j}^{n+1/2} \left|\phi_{j-1}^{n} - \phi_{j-1}^{n-1}\right| \\ &+ \left|\lambda\phi_{j}^{n-1}\left(g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2}\right)\right|. \end{aligned}$$

$$(2.26)$$

Summing over j and cancelling telescoping terms we obtain

$$\begin{split} &\sum_{j=1}^{M} \left| \phi_{j}^{n+1/2} - \phi_{j}^{n-1/2} \right| \\ &\leqslant \sum_{j=1}^{M} \left| \phi_{j}^{n} - \phi_{j}^{n-1} \right| + \lambda g_{V,1}^{n+1/2} |\phi_{0}^{n} - \phi_{0}^{n-1}| + \lambda \phi_{M}^{n-1} \left| g_{V,M+1}^{n+1/2} - g_{V,M+1}^{n-1/2} \right| \\ &- \lambda g_{V,M+1}^{n+1/2} \left| \phi_{M}^{n} - \phi_{M}^{n-1} \right|. \end{split}$$

The last inequality implies

$$\sum_{j=1}^{M} \left| \phi_{j}^{n+1/2} - \phi_{j}^{n-1/2} \right| \leq \sum_{j=1}^{M} \left| \phi_{j}^{n} - \phi_{j}^{n-1} \right| + \left| r^{n} - r^{n-1} \right| + \left| \mathcal{G}(t^{n}) - \mathcal{G}(t^{n-1}) \right|.$$

We observe that

$$\phi_{j}^{n} - \phi_{j}^{n-1} = \left(1 - B_{j+1/2}^{n-1} - A_{j-1/2}^{n-1}\right) \left(\phi_{j}^{n-1/2} - \phi_{j}^{n-3/2}\right) + A_{j+1/2}^{n-1} \left(\phi_{j+1}^{n-1/2} - \phi_{j+1}^{n-3/2}\right) + B_{j-1/2}^{n-1} \left(\phi_{j-1}^{n-1/2} - \phi_{j-1}^{n-3/2}\right),$$
(2.27)

where

$$\begin{split} A_{j+1/2}^{n-1} &= -\lambda \int_0^1 \partial_1 \varphi \big(\theta \phi_{j+1}^{n-1/2} + (1-\theta) \theta \phi_{j+1}^{n-3/2}, \theta \phi_{j+1}^{n-1/2} + (1-\theta) \theta \phi_{j+1}^{n-3/2} \big) \, \mathrm{d}\theta, \\ B_{j+1/2}^{n-1} &= \lambda \int_0^1 \partial_2 \varphi \big(\theta \phi_{j+1}^{n-1/2} + (1-\theta) \theta \phi_{j+1}^{n-3/2}, \theta \phi_{j+1}^{n-1/2} + (1-\theta) \theta \phi_{j+1}^{n-3/2} \big) \, \mathrm{d}\theta. \end{split}$$

Herein $\varphi(\phi_{j+1}, \phi_j) = \phi_j p_V(\phi_{j+1})$ and $\partial_i \varphi$ denotes the partial derivative of φ with respect to the *i*-th argument (i = 1, 2). Since $\phi, p_V(\phi) \ge 0$ and $p'_V(\phi) \le 0$, the function $\varphi(\phi_{j+1}, \phi_j)$ is nonincreasing with respect to ϕ_{j+1} and nondecreasing with respect to ϕ_j . This implies (together with the CFL condition)

$$0 \leqslant A_{j+1/2}^{n-1}, B_{j+1/2}^{n-1} \leqslant \frac{1}{2}.$$
(2.28)

The remainder of the proof is similar to the proof of Lemma 5.6 in [28]. Details are omitted. $\hfill \Box$

Now, we are ready to prove the convergence of ϕ^{Δ} as $\Delta \to \mathbf{0}$.

Lemma 2.7. The functions ϕ^{Δ} defined by (2.15) converge in $L^{1}(\Pi_{T})$ and boundedly *a.e.* a along subsequence to a limit function $\phi \in C([0,T], L^{1}(-L,L)) \cap L^{\infty}(\Pi_{T}))$.

Proof. The proof is a standard argument using the L^{∞} estimate (Lemma 2.2), the uniform spatial variation bound (Lemma 2.3), and the L^1 Lipschitz continuity in time estimate (Lemma 2.6).

In order to show that the limit function ϕ identified in Lemma 2.7 is a weak solution in the sense of Definition 2.1, we must also prove the convergence of the flux approximations. Instead of showing that the approximations $\{g_{V,j}^{n+1/2}\}$ converge we show that the approximations $\{h_j^{n+1/2}\}$ converge, where we define

$$h_j^{n+1/2} := \phi_j^{n+1/2} g_{V,j}^{n+1/2}$$
 for all $j = 1, \dots, M$ and $n = 0, \dots, \mathcal{N}$,

and extend these quantities to functions defined on Π_T by

$$h^{\Delta}(x,t) := \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \chi_j(x) \chi^n(t) h_j^{n+1/2}.$$

Now, we require additional time continuity estimates, which is the contents of the following lemma. Its proof is very similar to that of Lemmas 5.8 and 5.9 in [28], and is therefore omitted.

Lemma 2.8. The following uniform estimates hold for $n \ge 1$, where the constant Ω_4 is independent of Δ :

$$\sum_{j=1}^{M} \left| \phi_j^{n+1} - \phi_j^n \right| \leqslant \Omega_4, \qquad \Omega_4 := \Omega_2 + \mathrm{TV}(s) + \mathrm{TV}(r), \tag{2.29}$$

$$\sum_{j=1}^{M} \left| \phi_{j}^{n+1/2} - \phi_{j}^{n} \right| \leq \Omega_{1} + \Omega_{4}.$$
(2.30)

The following lemma is needed to establish a spatial variation bound on the approximations $h_j^{n+1/2}$.

Lemma 2.9. There exists a constant Ω_5 that is independent of Δ such that

$$\sum_{j=1}^{M} \phi_j^n \left| \Delta_+ g_{V,j}^{n+1/2} \right| \leqslant \Omega_5.$$

Proof. From the first half-step of the scheme we get

$$\phi_j^{n+1/2} - \phi_j^n = -\lambda \big(\phi_j^n \Delta_+ g_{V,j}^{n+1/2} + g_{V,j}^{n+1/2} \Delta_+ \phi_{j-1}^n \big),$$

which can be rerranged as

$$\lambda \phi_j^n \Delta_+ g_{V,j}^{n+1/2} = -\left(\phi_j^{n+1/2} - \phi_j^n\right) - \lambda g_{V,j}^{n+1/2} \Delta_+ \phi_{j-1}^n.$$

Taking absolute values and summing over $j = 1, \ldots, M$ we get

$$\lambda \sum_{j=1}^{M} \phi_{j}^{n} \left| \Delta_{+} g_{V,j}^{n+1/2} \right| \leq \sum_{j=1}^{M} \left| \phi_{j}^{n+1/2} - \phi_{j}^{n} \right| + \lambda \sum_{j=1}^{M} g_{V,j}^{n+1/2} \left| \Delta_{+} \phi_{j-1}^{n} \right|.$$

From Lemma 2.1 and the CFL condition (2.14) we have

$$\sum_{j=1}^{M} \phi_{j}^{n} \left| \Delta_{+} g_{V,j}^{n+1/2} \right| \leq \frac{1}{\lambda} \sum_{j=1}^{M} \left| \phi_{j}^{n+1/2} - \phi_{j}^{n} \right| + \sum_{j=1}^{M} \left| \Delta_{+} \phi_{j-1}^{n} \right|.$$

The result is obtained from (2.30) in Lemma 2.8 and Lemma 2.3.

We are now ready to bound the spatial variation of the approximations $h_j^{n+1/2}$. Lemma 2.10. There exists a constant Ω_6 that is independent of Δ such that

$$\sum_{j=1}^{M} |h_{j+1}^{n+1/2} - h_j^{n+1/2}| \le \Omega_6.$$
(2.31)

Proof. The first part of scheme (2.6) can be written as

$$\phi_j^{n+1/2} = \phi_j^n - \lambda \left(\phi_j^n g_{V,j+1}^{n+1/2} + g_{V,j}^{n+1/2} \left(\phi_j^{n+1/2} - \phi_{j-1}^n \right) \right) + \lambda \phi_j^{n+1/2} g_{V,j}^{n+1/2}.$$

Applying the spatial difference operator to the above equation we get

$$\Delta_{+}\phi_{j}^{n+1/2} = \left(1 - \lambda g_{V,j+1}^{n+1/2}\right)\Delta_{+}\phi_{j}^{n} + \lambda \Delta_{+}h_{j}^{n+1/2} - \lambda \phi_{j+1}^{n}\Delta_{+}g_{V,j+1}^{n+1/2} - \lambda \Delta_{+}g_{V,j}^{n+1/2}\left(\phi_{j}^{n+1/2} - \phi_{j}^{n}\right) + \lambda g_{V,j}^{n+1/2}\Delta_{+}\phi_{j-1}^{n} - \lambda g_{V,j+1}^{n+1/2}\Delta_{+}\phi_{j}^{n+1/2}$$

Thus

$$\begin{split} \lambda \Delta_{+} h_{j}^{n+1/2} &= \Delta_{+} \phi_{j}^{n+1/2} - \left(1 - \lambda g_{V,j+1}^{n+1/2}\right) \Delta_{+} \phi_{j}^{n} + \lambda \phi_{j+1}^{n} \Delta_{+} g_{V,j+1}^{n+1/2} \\ &+ \lambda \Delta_{+} g_{V,j}^{n+1/2} \left(\phi_{j}^{n+1/2} - \phi_{j}^{n}\right) + \lambda g_{V,j}^{n+1/2} \Delta_{+} \phi_{j-1}^{n} + \lambda g_{V,j+1}^{n+1/2} \Delta_{+} \phi_{j}^{n+1/2} \end{split}$$

After taking absolute values and using $|\Delta_+ g_{V,j}^{n+1/2}| \leq \alpha_V$, Lemma 2.1 and the CFL condition (2.14) we get

$$\begin{split} \lambda \left| \Delta_{+} h_{j}^{n+1/2} \right| \\ \leqslant 2 \left| \Delta_{+} \phi_{j}^{n+1/2} \right| + \left| \Delta_{+} \phi_{j}^{n} \right| + \lambda \phi_{j+1}^{n} \left| \Delta_{+} g_{V,j+1}^{n+1/2} \right| + \left| \phi_{j}^{n+1/2} - \phi_{j}^{n} \right| + \left| \Delta_{+} \phi_{j-1}^{n} \right| \\ \text{Summing over } j = 1, \dots, M \text{ we get} \end{split}$$

$$\sum_{j=1}^{M} \left| \Delta_{+} h_{j}^{n+1/2} \right|$$

MULTICLASS TRAFFIC WITH DISCONTINUOUS VELOCITY

$$\leq \frac{2}{\lambda} \sum_{j=1}^{M} \left| \Delta_{+} \phi_{j}^{n+1/2} \right| + \frac{2}{\lambda} \sum_{j=1}^{M} \left| \Delta_{+} \phi_{j}^{n} \right| + \frac{1}{\lambda} \sum_{j=1}^{M} \left| \phi_{j}^{n+1/2} - \phi_{j}^{n} \right| + \sum_{j=1}^{M} \phi_{j+1}^{n} \left| \Delta_{+} g_{V,j+1}^{n+1/2} \right|.$$

Finally, the result follows from Lemma 2.3, (2.30) in Lemma 2.8, and Lemma 2.9. $\hfill \Box$

The following lemma is required to prove the L^1 Lipschitz continuity in time and spatial variation bounds on $\{h_j^{n+1/2}\}.$

Lemma 2.11. There exists a constant Ω_7 that is independent of Δ such that

$$\Delta x \sum_{j=1}^{M} \sum_{n=1}^{N} \phi_j^{n-1} \left| g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2} \right| \leq \Omega_7.$$
(2.32)

Proof. From (2.26) we get

$$\begin{split} \lambda \phi_{j}^{n-1} \left| g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2} \right| &\leqslant \left(1 - \lambda g_{V,j+2}^{n+1/2} \right) \left| \phi_{j+1}^{n} - \phi_{j+1}^{n-1} \right| + \lambda g_{V,j+1}^{n+1/2} \left| \phi_{j}^{n} - \phi_{j}^{n-1} \right| \\ &- \left| \phi_{j+1}^{n+1/2} - \phi_{j+1}^{n-1/2} \right| + \lambda \phi_{j+1}^{n-1} \left| g_{V,j+2}^{n+1/2} - g_{V,j+2}^{n-1/2} \right|. \end{split}$$

By induction we obtain

$$\lambda \phi_{j}^{n-1} \left| g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2} \right| \leq \sum_{k=j+1}^{M} \left| \phi_{k}^{n} - \phi_{k}^{n-1} \right| + \left| g_{V,M+1}^{n+1/2} - g_{V,M+1}^{n-1/2} \right| - \sum_{k=j+1}^{M} \left| \phi_{k}^{n+1/2} - \phi_{k}^{n-1/2} \right| + \left| \phi_{j}^{n} - \phi_{j}^{n-1} \right|.$$

$$(2.33)$$

Recalling (2.27) we have

$$\begin{split} &\sum_{k=j+1}^{M} \left| \phi_{k}^{n} - \phi_{k}^{n-1} \right| \\ &\leqslant \sum_{k=j+1}^{M} \left(1 - B_{k+1/2}^{n-1} - A_{k-1/2}^{n-1} \right) \left| \phi_{k}^{n-1/2} - \phi_{k}^{n-3/2} \right| \\ &+ \sum_{k=j+1}^{M} A_{k+1/2}^{n-1} \left| \phi_{k+1}^{n-1/2} - \phi_{k+1}^{n-3/2} \right| + \sum_{k=j+1}^{M} B_{k-1/2}^{n-1/2} \left| \phi_{k-1}^{n-1/2} - \phi_{k-1}^{n-3/2} \right|. \end{split}$$

Cancelling telescoping terms and applying (2.28) yields

$$\sum_{k=j+1}^{M} \left| \phi_k^n - \phi_k^{n-1} \right|$$

$$\leq \sum_{k=j+1}^{M} \left| \phi_k^{n-1/2} - \phi_k^{n-3/2} \right| + \frac{1}{2} \left| \phi_{M+1}^{n-1/2} - \phi_{M+1}^{n-3/2} \right| + \frac{1}{2} \left| \phi_j^{n-1/2} - \phi_j^{n-3/2} \right|.$$

Then (2.33) becomes

$$\begin{split} \lambda \phi_{j}^{n-1} \big| g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2} \big| \\ \leqslant \sum_{k=j+1}^{M} \big| \phi_{k}^{n-1/2} - \phi_{k}^{n-3/2} \big| - \sum_{k=j+1}^{M} \big| \phi_{k}^{n+1/2} - \phi_{k}^{n-1/2} \big| + \frac{1}{2} \big| \phi_{M+1}^{n-1/2} - \phi_{M+1}^{n-3/2} \big| \end{split}$$

$$+\frac{1}{2}\left|\phi_{j}^{n-1/2}-\phi_{j}^{n-3/2}\right|+\left|\phi_{j}^{n}-\phi_{j}^{n-1}\right|+\left|g_{V,M+1}^{n+1/2}-g_{V,M+1}^{n-1/2}\right|$$

Summing over $n \ge 2$ and j = 1, ..., M, cancelling telescoping terms and multiplying the result by Δx we get

$$\Delta x \sum_{j=1}^{M} \sum_{n=2}^{N} \phi_j^{n-1} \left| g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2} \right| \leqslant S_1 + \dots + S_5,$$

where we define

$$S_{1} := \frac{2L}{\lambda} \sum_{j=1}^{M} |\phi_{j}^{3/2} - \phi_{j}^{1/2}|, \quad S_{2} := \frac{L}{\lambda} \sum_{n=2}^{N} |s^{n-1/2} - s^{n-3/2}|,$$

$$S_{3} := \frac{\Delta x}{\lambda} \sum_{j=1}^{M} \sum_{n=2}^{N} |\phi_{j}^{n} - \phi_{j}^{n-1}|, \quad S_{4} := \frac{2L}{\lambda} \sum_{n=2}^{N} |g_{V,M+1}^{n+1/2} - g_{V,M+1}^{n-1/2}|,$$

$$S_{5} := \frac{\Delta x}{2\lambda} \sum_{j=1}^{M} \sum_{n=2}^{N} |\phi_{j}^{n-1/2} - \phi_{j}^{n-3/2}|.$$

In view of the bounds established so far, there holds

$$S_1 \leqslant \frac{2L}{\lambda}\Omega_3, \quad S_2 \leqslant \frac{L}{\lambda}\mathrm{TV}(s), \quad S_3 \leqslant \Omega_4 T, \quad S_4 \leqslant \frac{2L}{\lambda}\mathrm{TV}(\mathcal{G}), \quad S_5 \leqslant \frac{\Omega_3}{2}T.$$

These bounds in conjunction with $|g_{V,j}^{3/2} - g_{V,j}^{1/2}| \leq \alpha_V$ imply that there exists a constant Ω_7 such that (2.32) is valid.

Lemma 2.12. There exists a constant Ω_8 that is independent of Δ such that

$$\Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \left| h_{j+1}^{n+1/2} - h_{j}^{n+1/2} \right| + \Delta x \sum_{n=1}^{\mathcal{N}} \sum_{j=1}^{M} \left| h_{j}^{n+1/2} - h_{j}^{n-1/2} \right| \leq \Omega_{8}.$$
(2.34)

Proof. In light of the spatial variation bound (2.31) we find that

$$\Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \left| h_{j+1}^{n+1/2} - h_{j}^{n+1/2} \right| \leqslant \Omega_{6} T.$$

The first part of (2.6) implies

$$\begin{split} \phi_{j}^{n+1/2} &- \phi_{j}^{n-1/2} \\ = \left(1 - \lambda g_{V,j+1}^{n+1/2}\right) \left(\phi_{j}^{n} - \phi_{j}^{n-1}\right) + \lambda \left(h_{j}^{n+1/2} - h_{j}^{n-1/2}\right) - \lambda g_{V,j}^{n+1/2} \left(\phi_{j}^{n+1/2} - \phi_{j-1}^{n}\right) \\ &+ \lambda g_{V,j}^{n-1/2} \left(\phi_{j}^{n-1/2} - \phi_{j-1}^{n-1}\right) - \lambda \phi_{j}^{n-1} \left(g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2}\right) \\ = \left(1 - \lambda g_{V,j+1}^{n+1/2}\right) \left(\phi_{j}^{n} - \phi_{j}^{n-1}\right) + \lambda \left(h_{j}^{n+1/2} - h_{j}^{n-1/2}\right) - \lambda g_{V,j}^{n+1/2} \left(\phi_{j}^{n+1/2} - \phi_{j}^{n}\right) \\ &- \lambda g_{V,j}^{n+1/2} \Delta_{+} \phi_{j-1}^{n} + \lambda g_{V,j}^{n-1/2} \Delta_{+} \phi_{j-1}^{n-1} + \lambda g_{V,j}^{n-1/2} \left(\phi_{j}^{n-1/2} - \phi_{j}^{n-1}\right) \\ &- \lambda \phi_{j}^{n-1} \left(g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2}\right). \end{split}$$

Consequently,

$$\begin{split} \lambda \big(h_j^{n+1/2} - h_j^{n-1/2} \big) \\ &= \big(\phi_j^{n+1/2} - \phi_j^{n-1/2} \big) - \big(1 - \lambda g_{V,j+1}^{n+1/2} \big) \big(\phi_j^n - \phi_j^{n-1} \big) + \lambda g_{V,j}^{n+1/2} \big(\phi_j^{n+1/2} - \phi_j^n \big) \\ &+ \lambda g_{V,j}^{n+1/2} \Delta_+ \phi_{j-1}^n - \lambda g_{V,j}^{n-1/2} \Delta_+ \phi_{j-1}^{n-1} - \lambda g_{V,j}^{n-1/2} \big(\phi_j^{n-1/2} - \phi_j^{n-1} \big) \end{split}$$

$$+ \lambda \phi_j^{n-1} (g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2}).$$

Taking absolute values and using the CFL condition (2.14) we get

$$\begin{split} \lambda \left| h_{j}^{n+1/2} - h_{j}^{n-1/2} \right| \\ &\leqslant \left| \phi_{j}^{n+1/2} - \phi_{j}^{n-1/2} \right| + \left| \phi_{j}^{n} - \phi_{j}^{n-1} \right| + \left| \Delta_{+} \phi_{j-1}^{n} \right| + \left| \Delta_{+} \phi_{j-1}^{n-1} \right| \\ &+ \left| \phi_{j}^{n+1/2} - \phi_{j}^{n} \right| + \left| \phi_{j}^{n-1/2} - \phi_{j}^{n-1} \right| + \lambda \phi_{j}^{n-1} \left| g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2} \right|. \end{split}$$

Multiplying this inequality by Δx and summing over j and n we get

$$\Delta x \sum_{n=1}^{N} \sum_{j=1}^{M} \left| h_j^{n+1/2} - h_j^{n-1/2} \right| \leq U_1 + \dots + U_5,$$

where we define

$$U_{1} := \frac{\Delta x}{\lambda} \sum_{n=1}^{N} \sum_{j=1}^{M} |\phi_{j}^{n+1/2} - \phi_{j}^{n-1/2}|, \quad U_{2} := \frac{\Delta x}{\lambda} \sum_{n=1}^{N} \sum_{j=1}^{M} |\phi_{j}^{n} - \phi_{j}^{n-1}|,$$
$$U_{3} := \frac{2\Delta x}{\lambda} \sum_{n=1}^{N} \sum_{j=1}^{M} |\Delta_{+}\phi_{j-1}^{n}|, \quad U_{4} := \frac{2\Delta x}{\lambda} \sum_{n=1}^{N} \sum_{j=1}^{M} |\phi_{j}^{n+1/2} - \phi_{j}^{n}|,$$
$$U_{5} := \Delta x \sum_{n=1}^{N} \sum_{j=1}^{M} \phi_{j}^{n-1} |g_{V,j+1}^{n+1/2} - g_{V,j+1}^{n-1/2}|.$$

From (2.30), (2.29), (2.18), (2.30) and (2.32) we have

$$U_1 \leq \Omega_3 T, \quad U_2 \leq \Omega_4 T, \quad U_3 \leq 2 \big(\mathrm{TV}(\phi_0) + \mathrm{TV}(s) + \mathrm{TV}(r) \big) T,$$

$$U_4 \leq 2 \big(\Omega_1 + \Omega_4 \big) T, \quad U_5 \leq \Omega_7.$$

Combining these bounds we see that there exists a constant Ω_8 that is independent of Δ such that (2.34) is valid.

Lemma 2.13. The functions h^{Δ} converge in $L^1(\Pi_T)$ and boundedly a.e. along subsequence to some limit function $w \in L^1(-L,L) \cap L^{\infty}(-L,L)$. Moreover, by a suitable choice of a subsequence, we have $w(x,t) \in \tilde{Q}(\phi(x,t))$ a.e. in Π_T , where $\phi(x,t)$ is the limit of Lemma 2.7.

Proof. We observe that $|h_j^{n+1/2}| \leq \phi_{\max} \alpha_V$. Then by Helly's theorem [23] there exists a function $w \in L^1(\Pi_T)$ such that $h^{\Delta} \to w$ along a subsequence in $L^1(\Pi_T)$ and boundedly a.e. in Π_T . To prove the second assertion, we define $Q(\phi) := \phi g_V(\phi)$ and \tilde{Q} denote the multivalued version of Q. Assume (by extracting further subsequences if necessary) that $\phi^{\Delta} \to \phi$, $h^{\Delta} \to w$ in $L^1(\Pi_T)$ and fix a point $(x,t) \in \Pi_T$ where $\phi^{\Delta}(x,t) \to \phi(x,t)$ and $h^{\Delta}(x,t) \to w(x,t)$ as $\Delta \to \mathbf{0}$. First, we consider the case $\phi(x,t) = \phi^*$. Lemma 2.1 implies that $0 \leq h^{\Delta}(x,t) \leq \alpha_V \phi^{\Delta}(x,t)$. Then passing to the limit in the above inequality we obtain

$$w(x,t) \in [0, \alpha_V \phi^*] = \tilde{Q}(\phi^*).$$

In case $\phi(x,t) \neq \phi^*$ first we consider $\phi(x,t) < \phi^*$, then $\tilde{Q}(\phi(x,t)) = \alpha_V \phi(x,t)$. For sufficiently small Δ the inequality $\phi^{\Delta}(x,t) < \phi^*$ implies that $\phi_j^{n+1/2} < \phi^*$ and $\tilde{g}_V(\phi_j^{n+1/2}) = \{\alpha_V\}$. Then, by Lemma 2.1 we get

$$h^{\Delta}(x,t) = \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \chi_j(x) \chi^n(t) h_j^{n+1/2} = \alpha_V \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \chi_j(x) \chi^n(t) \phi_j^{n+1/2} = \alpha_V \phi^{\Delta}(x,t).$$

Thus $w(x,t) = \lim h^{\Delta}(x,t) = \alpha_V \lim \phi^{\Delta}(x,t) = \alpha_V \phi(x,t) = \tilde{Q}(\phi(x,t)).$

In the case $\phi(x,t) > \phi^*$ there holds $\tilde{Q}(\phi(x,t)) = 0$. In this case it is necessary extend $\{g_{V,j}^{n+1/2}\}$ to functions defined on Π_T by

$$g_{V}^{\Delta}(x,t) = \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \chi_{j}(x) \chi^{n}(t) g_{V,j}^{n+1/2},$$

and we need to utilize the following consequence of Lemma 2.1:

$$g_V^{\Delta}(x,t) \in \tilde{g}_V(\phi^{\Delta}(x,t)), \quad (x,t) \in \Pi_T.$$

For sufficiently small Δ , $\phi^{\Delta}(x,t) > \phi^*$ implies that $\tilde{g}_V(\phi^{\Delta}(x,t)) = \{0\}$, hence $g_V^{\Delta}(x,t) = 0$. Finally observe that $0 \leq h^{\Delta}(x,t) \leq \phi^{\Delta}(x,t) \cdot g_V^{\Delta}(x,t) = 0$ for sufficiently small Δ . Hence $w(x,t) = \tilde{Q}(\phi(x,t)) = 0$.

Theorem 2.14 (Main result). The functions ϕ^{Δ} converge in $L^{1}(\Pi_{T})$ and boundedly a.e. along subsequence to some $\phi \in C([0,T], L^{1}(-L,L)) \cap L^{\infty}(\Pi_{T})$. The limit function $\phi(x,t)$ is a weak solution in sense of Definition 2.1.

Proof. The convergence is ensured by Lemma 2.7. It remains to prove that the limit ϕ is a weak solution. Let us fix a point $(x,t) \in \Pi_T$, then Lemma 2.13 implies that $w(x,t) \in \tilde{Q}(\phi(x,t))$ a.e. in Π_T . If $\phi(x,t) \neq \phi^*$, then $\tilde{Q}(\phi(x,t)) = Q(\phi(x,t))$. Thus $w(x,t) = Q(\phi(x,t))$, then we define $q(x,t) = \phi_V(\phi) + Q(\phi(x,t)) = f(\phi(x,t))$. In the case where $\phi(x,t) = \phi^*$ we take $w(x,t) \in [0, \alpha_V \phi^*]$ and define

$$q(x,t) = \phi^* p_V(\phi^*) + w(x,t) \in [\phi^* p_V(\phi^*), \phi^* p_V(\phi^*) + \alpha_V \phi^*] = \tilde{f}(\phi^*).$$

In either case $q(x,t) \in \tilde{f}(\phi(x,t))$.

We note that the two steps of (2.6) imply

$$\phi_j^{n+1} - \phi_j^n + \lambda \left(\phi_j^n g_{V,j+1}^{n+1/2} - \phi_{j-1}^n g_{V,j}^{n+1/2} + \phi_j^{n+1/2} p_{V,j+1}^{n+1/2} - \phi_{j-1}^{n+1/2} p_{V,j}^{n+1/2} \right) = 0. \quad (2.35)$$

We now choose a test function $\psi \in C_0^1((-L, L) \times [0, T))$ and define $\psi_j^n := \psi(x_j, t^n)$. Multiplying (2.35) by $\Delta x \psi_j^n$ and summing the result over j and n yields

$$\begin{split} \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\phi_{j}^{n+1} - \phi_{j}^{n}}{\Delta t} \psi_{j}^{n} + \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\phi_{j}^{n} g_{V,j+1}^{n+1/2} - \phi_{j-1}^{n} g_{V,j}^{n+1/2}}{\Delta x} \psi_{j}^{n} \\ + \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\phi_{j}^{n+1/2} p_{V,j+1}^{n+1/2} - \phi_{j-1}^{n+1/2} p_{V,j}^{n+1/2}}{\Delta x} \psi_{j}^{n} = 0. \end{split}$$

A summation by parts yields

$$\Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \phi_{j}^{n+1} \frac{\psi_{j}^{n+1} - \psi_{j}^{n}}{\Delta t} + \Delta x \sum_{j=1}^{M} \phi_{j}^{0} \psi_{j}^{0} + \Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n} g_{V,j+1}^{n+1/2} + \Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n+1/2} p_{V,j+1}^{n+1/2} = 0.$$

$$(2.36)$$

An application of (2.12) yields, as $\Delta x, \Delta t \to 0$,

$$\Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \phi_j^{n+1} \frac{\psi_j^{n+1} - \psi_j^n}{\Delta t} = \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \phi_j^{n+1/2} \frac{\psi_j^{n+1} - \psi_j^n}{\Delta t} + \mathcal{O}(\Delta x).$$

This equation and Lemma 2.3 imply that the two first sums in (2.36) converge to

$$\int_{0}^{T} \int_{-L}^{L} \phi \psi_{t} \, \mathrm{d}x + \int_{-L}^{L} \phi_{0}(x) \psi(x,0) \, \mathrm{d}x \, \mathrm{d}t$$

Concerning the last term in (2.36), we get

$$\begin{split} \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n+1/2} p_{V,j+1}^{n+1/2} \\ &= \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n+1/2} p_{V,j}^{n+1/2} \\ &+ \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \left(\phi_{j}^{n+1/2} p_{V,j+1}^{n+1/2} - \phi_{j}^{n+1/2} p_{V,j}^{n+1/2} \right). \end{split}$$

By properties of the function p_V we get the estimate

 $\phi_j^{n+1/2} p_{V,j+1}^{n+1/2} - \phi_j^{n+1/2} p_{V,j}^{n+1/2} = \phi_j^{n+1/2} \left(p_{V,j+1}^{n+1/2} - p_{V,j}^{n+1/2} \right) \leqslant \phi_{\max} \| p_V' \|_{\infty} \Delta x.$ Thus

$$\left| \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\left(\psi_{j+1}^{n} - \psi_{j}^{n}\right)}{\Delta x} \left(\phi_{j}^{n+1/2} p_{V,j+1}^{n+1/2} - \phi_{j}^{n+1/2} p_{V,j}^{n+1/2}\right) \right| \\ \leqslant 2MT \phi_{\max} \Delta x \|\partial_{t} \psi\|_{\infty} \|p_{V}'\|_{\infty},$$

which goes to zero as $\Delta x \to 0$. Therefore the last term in (2.36) converges to

$$\int_0^T \int_{-L}^L \phi p_V(\phi) \psi_x \, \mathrm{d}x \, \mathrm{d}t$$

as $\Delta x \to 0$. The second term in (2.36) can be written as

$$\begin{aligned} \Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n} g_{V,j+1}^{n+1/2} \\ &= \Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n+1/2} g_{V,j}^{n+1/2} + \Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n} \Delta_{+} g_{V,j}^{n+1/2} \\ &+ \Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} g_{V,j}^{n+1/2} (\phi_{j}^{n} - \phi_{j}^{n+1/2}). \end{aligned}$$

Using Lemmas 2.9, 2.1, and 2.8 we get

$$\left| \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n} \Delta_{+} g_{V,j}^{n+1/2} \right| \leq \Delta x \|\partial_{x}\psi\|_{\infty} \Omega_{5}T,$$
$$\left| \Delta x \Delta t \sum_{n=0}^{\mathcal{N}} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} g_{V,j}^{n+1/2} (\phi_{j}^{n} - \phi_{j}^{n+1/2}) \right| \leq \alpha_{V} \Delta x \|\partial_{x}\psi\|_{\infty} (\Omega_{1} + \Omega_{4})T.$$

Consequently, as $\Delta x \to 0$,

$$\Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} \phi_{j}^{n} \Delta_{+} g_{V,j}^{n+1/2} \to 0,$$

$$\Delta x \Delta t \sum_{n=0}^{N} \sum_{j=1}^{M} \frac{\psi_{j+1}^{n} - \psi_{j}^{n}}{\Delta x} g_{V,j}^{n+1/2} (\phi_{j}^{n} - \phi_{j}^{n+1/2}) \to 0.$$

Then substituting $g_V^{\Delta}(x_j, t^n) = g_{V,j}^{n+1/2}$ and applying the dominated convergence theorem we obtain that the second term in (2.36) converges to

$$\int_0^T \int_{-L}^L w\psi_x \,\mathrm{d}x \,\mathrm{d}t.$$

Collecting the previous results we get

$$\int_0^T \int_{-L}^L (\phi \psi_t + q \psi_x) \, \mathrm{d}x \, \mathrm{d}t + \int_{-L}^L \phi_0(x) \psi(x,0) \, \mathrm{d}x = 0,$$

so ϕ is a weak solution in sense of Definition 2.1.

3. Extension to the MCLWR model. Algorithm 2.1 cannot be applied directly in a component-wise manner for each class *i* in the multiclass case (1.1)–(1.4), but we can first solve for the total density ϕ and then update the densities ϕ_1, \ldots, ϕ_N for each class. The multiclass version of the scalar scheme (1.6) can be written as

$$\phi_{i,j}^{n+1/2} = \phi_{i,j}^n - \lambda v_i^{\max} \left(\phi_{i,j}^n g_V \left(\phi_{j+1}^{n+1/2} \right) - \phi_{i,j-1}^n g_V \left(\phi_j^{n+1/2} \right) \right),$$

$$\phi_{i,j}^{n+1} = \phi_{i,j}^{n+1/2} - \lambda v_i^{\max} \left(\phi_{i,j}^{n+1/2} p_V \left(\phi_{j+1}^{n+1/2} \right) - \phi_{i,j-1}^{n+1/2} p_V \left(\phi_j^{n+1/2} \right) \right),$$

$$i = 1, \dots, N,$$
(3.1)

where the following quantity is an approximate value of the total density ϕ :

$$\phi_j^{n+1/2} := \phi_{1,j}^{n+1/2} + \dots + \phi_{N,j}^{n+1/2}.$$

In order to solve (3.1), we need to impose the non-standard boundary condition (1.4b). Recalling that $V(\phi) = g_V(\phi) + p_V(\phi)$ we can equivalently specify for the multiclass case the condition (2.2). The correspondence when $s(t) = \phi^*$ is

$$\mathcal{F}(t) = (\mathbf{v}^{\max})^{\mathrm{T}} \mathbf{s}(t) V(\phi^* -) \Leftrightarrow \mathcal{G}(t) = \alpha_V,$$

$$\mathcal{F}(t) = (\mathbf{v}^{\max})^{\mathrm{T}} \mathbf{s}(t) V(\phi^* +) \Leftrightarrow \mathcal{G}(t) = 0.$$
(3.2)

Coming back to (3.1), we define $\Phi_j^n := (\phi_{1,j}^n, \dots, \phi_{N,j}^n)^{\mathrm{T}}$. Summing over $i = 1, \dots, N$, assuming that g_V is evaluated at the new time step, and replacing $g_V(\phi_{j+1}^{n+1/2})$ by $g_{V,j+1}^{n+1/2}$, we get

$$\phi_j^{n+1/2} = \phi_j^n - \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \left(g_{V,j+1}^{n+1/2} \Phi_j^n - g_V(\phi_j^{n+1/2}) \Phi_{j-1}^n \right).$$
(3.3)

This can be rearranged as

$$\phi_j^{n+1/2} - \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_{j-1}^n g_V(\phi_j^{n+1/2}) = \phi_j^n - \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_j^n g_{V,j+1}^{n+1/2}.$$
 (3.4)

Let us now define the function

$$G_V(z;\Phi) := z - \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi g_V(z)$$

and denote by $\tilde{G}_V(\cdot; \Phi)$ its multivalued version (with respect to z). Then \tilde{G} is strictly increasing and has a unique inverse $z \mapsto \tilde{G}_V^{-1}(z; \Phi)$. Expressing (3.4) as

$$\tilde{G}_{V}\left(\phi_{j}^{n+1/2}; \Phi_{j-1}^{n}\right) = \phi_{j}^{n} - \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_{j}^{n} g_{V,j+1}^{n+1/2}$$
(3.5)

20

which allows us to obtain $\phi_j^{n+1/2}$ by applying $\tilde{G}_V^{-1}(z; \Phi)$ to both sides, that is $\int_{a}^{a+1/2} - \tilde{C}^{-1}(A^n - M(z)^{\max}) T \Phi^n z^{n+1/2} \cdot \Phi^n$

$$\prod_{j=1}^{n+1/2} = G_V^{-1} (\phi_j^n - \lambda(\boldsymbol{v}^{\max}))^T \Phi_j^n g_{V,j+1}^{n+1/2}; \Phi_{j-1}^n).$$

Now that $\phi_j^{n+1/2}$ is available, we solve for $g_{V,j}^{n+1/2}$ the equation

$$\phi_j^{n+1/2} = \phi_j^n - \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \left(g_{V,j+1}^{n+1/2} \Phi_j^n - g_{V,j}^{n+1/2} \Phi_{j-1}^n \right)$$

(cf. (3.3)). This yields

$$g_{V,j}^{n+1/2} = \frac{\phi_j^{n+1/2} - \phi_j^n + \lambda g_{V,j+1}^{n+1/2} (\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_j^n}{\lambda (\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_{j-1}^n},$$

provided that $\Phi_{j-1}^n \neq \mathbf{0}$. If $\Phi_{j-1}^n = \mathbf{0}$ then we set $g_{V,j}^{n+1/2} = g_V(\phi_j^{n+1/2})$. The numerical scheme for the multiclass model can be summarized in the following algorithm.

Algorithm 3.1 (BCOV scheme, multiclass case).

 $\begin{array}{ll} Input: \ approximate \ solution \ vector \ \{\phi_{i,j}^n\}_{j=1}^M, \ i=1,\ldots,N \ for \ t=t^n \\ g_{V,M+1}^{n+1/2} \leftarrow \mathcal{G}(\phi_{M+1}^{n+1/2}) \ (\text{using } (2.3) \ \text{and } (3.2)) \\ \textbf{do } j=M, M-1,\ldots,1 \\ \phi_j^{n+1/2} \leftarrow \tilde{G}_V^{-1}(\phi_j^n - \lambda g_{V,j+1}^{n+1/2}(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_j^n; \Phi_{j-1}^n) \\ \textbf{if } \Phi_{j-1}^n \neq \textbf{0 then} \\ g_{V,j}^{n+1/2} \leftarrow \frac{\phi_j^{n+1/2} - \phi_j^n + \lambda g_{V,j+1}^{n+1/2}(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_j^n}{\lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_{j-1}^n} \\ \textbf{else} \\ g_{V,j}^{n+1/2} \leftarrow g_V(\phi_j^{n+1/2}) \\ \textbf{endif} \\ \textbf{enddo} \\ \textbf{do } j=1,\ldots,N \\ \phi_{i,j}^{n+1/2} \leftarrow \phi_{i,j}^n - \lambda v_i^{\max}(\phi_{i,j}^n g_{V,j+1}^{n+1/2} - \phi_{i,j-1}^n g_{V,j}^{n+1/2}) \\ \textbf{enddo} \\ \textbf{enddo} \\ \textbf{do } j=1,\ldots,N \\ \phi_{i,j}^{n+1/2} \leftarrow \phi_{i,j}^n - \lambda v_i^{\max}(\phi_{i,j}^{n+1/2} p_V(\phi_{j+1}^{n+1/2}) - \phi_{i,j-1}^{n+1/2} p_V(\phi_j^{n+1/2})) \\ \textbf{enddo} \\ \textbf{enddo} \\ \textbf{do } i=1,\ldots,N \\ \phi_{i,j}^{n+1} \leftarrow \phi_{i,j}^{n+1/2} - \lambda v_i^{\max}(\phi_{i,j}^{n+1/2} p_V(\phi_{j+1}^{n+1/2}) - \phi_{i,j-1}^{n+1/2} p_V(\phi_j^{n+1/2})) \\ \textbf{enddo} \\ \textbf{enddo} \\ \textbf{output: approximate solution vectors } \{\phi_{i,j}^{n+1}\}_{j=1}^M, i=1,\ldots,N \ for \ t=t^{n+1} = t^n + \Delta t \end{aligned}$

Remark 3.1. We recall that the boundary condition $g_{V,M+1}^{n+1/2} = \mathcal{G}(\phi_{M+1/2}^{n+1/2})$ that appears in Algorithm 3.1 is defined using (2.3) for the total density $\phi_{M+1}^{n+1/2}$. We illustrate this boundary condition in Section 4.5.

The problem of interest to us is to show that \mathcal{D} is an invariant region of the scheme. To this end we first consider the evolution of the total density ϕ . Summing over $i = 1, \ldots, N$ the second equation in (3.1) yields

$$\phi_j^{n+1} = \phi_j^{n+1/2} - \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \left(p_V \left(\phi_{j+1}^{n+1/2} \right)^{n+1/2} \Phi_j^{n+1/2} - p_V \left(\phi_j^{n+1/2} \right) \Phi_{j-1}^n \right)$$

The above equation can be written in incremental form as

$$\phi_j^{n+1} = \phi_j^{n+1/2} + C_{j+1/2}^{n+1/2} \Delta_+ \phi_j^{n+1/2} - D_{j-1/2}^{n+1/2} \Delta_- \phi_j^{n+1/2}, \qquad (3.6)$$

where we define

$$\begin{split} C_{j+1/2}^{n+1/2} &:= \begin{cases} \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_{j}^{n+1/2} \frac{p_{V}(\phi_{j}^{n+1/2}) - p_{V}(\phi_{j+1}^{n+1/2})}{\phi_{j+1}^{n+1/2} - \phi_{j}^{n+1/2}} & \text{if } \phi_{j+1}^{n+1/2} \neq \phi_{j}^{n+1/2}, \\ 0 & \text{if } \phi_{j+1}^{n+1/2} = \phi_{j}^{n+1/2}, \end{cases} \\ D_{j-1/2}^{n+1/2} &:= \begin{cases} \lambda p_{V}(\phi_{j}^{n+1/2}) \frac{(\boldsymbol{v}^{\max})^{\mathrm{T}}(\Phi_{j}^{n+1/2} - \Phi_{j-1}^{n+1/2})}{\phi_{j}^{n+1/2} - \phi_{j-1}^{n+1/2}} & \text{if } \phi_{j}^{n+1/2} \neq \phi_{j-1}^{n+1/2}, \\ 0 & \text{if } \phi_{j}^{n+1/2} = \phi_{j-1}^{n+1/2}, \end{cases} \\ \end{split}$$

Since $p_V(\phi)$ is a non-increasing positive function we have $C_{j+1/2}^{n+1/2}, D_{j-1/2}^{n+1/2} \ge 0$. To ensure that $|C_{j+1/2}^{n+1/2}| \le 1/2$ and $|D_{j-1/2}^{n+1/2}| \le 1/2$ we impose the CFL condition

$$\lambda \phi_{\max} \max_{1 \leqslant j \leqslant M} \left| p_V'(\phi_j^n) \right| \cdot \max_{1 \leqslant i \leqslant N} v_i^{\max} \leqslant \frac{1}{2}, \quad \lambda \max_{1 \leqslant j \leqslant M} p(\phi_j^n) \cdot \max_{1 \leqslant i \leqslant N} v_i^{\max} \leqslant \frac{1}{2}.$$
(3.7)

Lemma 3.1. Assume that

$$\Phi_j^0 \in \mathcal{D} \quad \text{for } j = 1, \dots, M. \tag{3.8}$$

Then $\Phi_j^n, \Phi_j^{n+1/2} \in \mathcal{D}$ for $j = 1, \dots, M$.

Proof. We claim that

$$\Phi_j^{n+1/2} \in \mathcal{D} \quad \text{for all } j = 1, \dots, M$$

$$\Rightarrow g_{V,j}^{n+1/2} \in [0, \alpha_V] \quad \text{for all } j = 1, \dots, M.$$
(3.9)

We consider first the case $\Phi_{j-1}^n = \mathbf{0}$. Then the result follows from the definition of the function $g_V(z)$ and (2.1). Let $\Phi_{j-1}^n \neq \mathbf{0}$. Summing over $i = 1, \ldots, N$ the first equation in (3.1) yields

$$\phi_j^{n+1/2} = \tilde{G}_V^{-1} \big(\phi_j^n - \lambda (\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_j^n g_{V,j+1}^{n+1/2}; \Phi_{j-1}^n \big).$$

Using (3.5) and (3.3) we find that

$$\phi_j^{n+1/2} - \lambda(\boldsymbol{v}^{\max})^{\mathrm{T}} \Phi_{j-1}^n g_{V,j}^{n+1/2} \in \tilde{G}_V(\phi_j^{n+1/2}; \Phi_{j-1}^n).$$

Thus, a straightforward case-by-case study and (3.5) prove that (3.9) is valid. The remainder of the proof is similar to the proof of Lemma 2.2.

4. Numerical examples. We now present some numerical simulations to illustrate the behaviour of solutions to system (1.1) by using Algorithms 2.1 and 3.1 for the scalar and multiclass case, respectively. In the scalar case, we compare numerical approximations with those generated by the scheme (1.7) proposed by Towers in [28]. To this end we choose the discontinuous velocity function

$$V(\phi) = \begin{cases} 1 - \phi/\phi_{\max} & \text{for } 0 \leqslant \phi \leqslant \phi^*, \\ -w_f(1 - \phi_{\max}/\phi) & \text{for } \phi^* < \phi \leqslant \phi_{\max} \end{cases}$$

where $\phi^* = 0.5$, $w_f = 0.2$, $\phi_{max} = 1$, and $\alpha_V = 0.3$.

In all numerical experiments computations are performed on a finite interval [-1,1] that is subdivided into M subintervals of length $\Delta x = 2/M$, and the time step is computed by $\Delta t = \Delta x/2$ in the scalar case (N = 1) and $\Delta t = \Delta x/(2 \max\{v_1^{\max}, \ldots, v_N^{\max}\})$ in the multiclass case $N \ge 2$. These choices ensure that the respective CFL conditions (2.14) and (3.7) are satisfied.



FIGURE 3. Example 1: numerical solution with M = 800 and comparison with the exact solution of the Riemann problem (a) with $\phi_{\rm L} = 0.3$ and $\phi_{\rm R} = 0.9$ at simulated time T = 1.8, (b) with $\phi_{\rm L} = 0.9$ and $\phi_{\rm R} = 0.3$ at simulated time T = 1.5. Here and in Figures 4 and 5 we label with 'Towers scheme' the scheme (1.7) proposed in [28] and by 'BCOV scheme' the scheme of Algorithm 2.1 advanced in the present work.

4.1. Example 1: scalar Riemann problem (N = 1). We consider the Riemann problem for the scalar equation $\partial_t \phi + \partial_x (\phi V(\phi)) = 0$ with initial data

$$\phi_0(x) = \begin{cases} \phi_{\rm L} & \text{for } x < 0.2, \\ \phi_{\rm R} & \text{for } x \ge 0.2 \end{cases}$$

$$\tag{4.1}$$

(no boundary conditions are involved). For $\phi_{\rm L} = 0.3$ and $\phi_{\rm R} = 0.9$, the solution consists of two shock waves with negative velocities of propagation, namely a shock wave connecting $\phi_{\rm L}$ with ϕ^* that travels velocity $\sigma_1 = -0.55$ and another shock wave connecting ϕ^* with $\phi_{\rm R}$ with velocity $\sigma_2 = -0.2$. Figure 3 (a) shows the numerical approximations to the solution of this problem computed with M = 800for both schemes at simulated time T = 1.8.

For $\phi_{\rm L} = 0.9$ and $\phi_{\rm R} = 0.3$, the solution consists of a shock wave connecting $\phi_{\rm L}$ with ϕ^* that travels at velocity $\sigma_1 = -0.575$ and a rarefaction wave connecting ϕ^* with $\phi_{\rm R}$. In Figure 3 (b) we display the numerical solutions computed with M = 800 for both schemes at simulated time T = 1.5. In both scenarios, all waves are approximated correctly by both schemes.

4.2. Example 2: scalar problem (N = 1), smooth initial datum. In this example we compare numerical approximations for equation (1.1) obtained by both schemes (Towers scheme (1.7) and Algorithm 2.1), starting from the initial function $\phi_0(x) = \exp(-(x + 0.2)^2/(0.04))$ for $x \in [-1, 1]$. Numerical approximations are computed at simulated times T = 0.1 and T = 0.3 with discretizations $M = 100 \times 2^l$, $l = 0, 1, \ldots, 4$. Table 1 displays the corresponding approximate L^1 errors obtained by utilizing a reference solution computed by the Towers scheme with $M_{\text{ref}} = 12800$, We observe that the approximate L^1 errors decrease as the grid is refined. In

R. BÜRGER, C. CHALONS, R. ORDOÑEZ, AND L. M. VILLADA

	T = 0.1		T = 0.3	
	Towers	BCOV	Towers	BCOV
M	$e_M(\phi^{\Delta})$	$e_M(\phi^{\Delta})$	$e_M(\phi^{\Delta})$	$e_M(\phi^{\Delta})$
100	1.32e-2	1.76e-2	1.63e-2	2.39e-2
200	6.55e-3	9.22e-3	8.59e-3	1.31e-2
400	3.29e-3	4.46e-3	4.25e-3	6.46e-3
800	1.72e-3	2.403 - 3	2.12e-3	3.31e-3
1600	8.00e-4	1.18e-3	9.29e-4	1.563-3

TABLE 1. Example 2: approximate L^1 errors $e_M(u)$ with $\Delta x = 2/M$.



FIGURE 4. Example 2: numerical solutions for M = 100 at simulated times (a) T = 0.1, (b) T = 0.3.

Figure 4 where we display the numerical approximations for M = 100 and compared with the reference solution.

4.3. Example 3: scalar problem (N = 1), non-standard boundary condition. This example comes from [28, Example 6.2] and is designed to illustrate that when $s(t^n) = \phi^*$, the solutions depend on the boundary condition $\mathcal{F}(t) \in \tilde{f}(\phi^*)$. For this example we consider the Riemann problem with initial data (4.1) with $\phi_{\rm L} = 1/4$ and $\phi_{\rm R} = 1/2$. We compute the solution twice, once using $\mathcal{G}(t) = \alpha_V$ (equivalently, $\mathcal{F}(t) = 1/2$), and the second time using $\mathcal{G}(t) = 0$ (equivalently, $\mathcal{F}(t) = 1/4$). As shown in Figure 5, in the first case the solution corresponds to a shock wave connecting $\phi_{\rm L}$ with $\phi_{\rm R}$ with speed of propagation $\sigma = 1$, and in the second case the solution corresponds to a stationary shock ($\sigma = 0$) connecting $\phi_{\rm L}$ with $\phi_{\rm R}$.

4.4. Example 4: multiclass case (N = 3), preservation of invariant region. To illustrate the invariant region property of the proposed scheme (Lemma 3.1), we consider the case N = 3 and the Riemann initial data

$$\phi_0(x) = \begin{cases} (0.1, 0.1, 0.1)^{\mathrm{T}} & \text{for } x < 0.5, \\ (0.4, 0.5, 0.1)^{\mathrm{T}} & \text{for } x \ge 0.5, \end{cases}$$



FIGURE 5. Example 3: numerical solutions depending on the boundary conditions $\mathcal{F}(t) \in \tilde{f}(\phi^*)$ with M = 1600 at simulated time T = 0.5, with (a) $\mathcal{F}(t) \in \tilde{f}(\phi^*-)$ (free flow), (b) $\mathcal{F}(t) \in \tilde{f}(\phi^*+)$ (congested flow).



FIGURE 6. Example 4: density profiles simulated with M = 1600 at (a) T = 0.2, (b) T = 0.4, (c) T = 0.6.

with velocities $\boldsymbol{v}^{\max} = (1,3,10)^{\mathrm{T}}$. The solution consists of a stationary shock plus two shock waves that travel with negative velocities. The numerical simulation at three simulated times is displayed in Figure 6. The profile for each class and the total density are displayed in this figure. Furthermore we can see that the profile of the total density in Figure 6 looks like the profile of Figure 3 (a).

4.5. Example 5: multiclass case (N = 3), non-standard boundary condition. It is the purpose of this example to illustrate the boundary condition

$$g_{V,M+1}^{n+1/2} = \mathcal{G}(\phi_{M+1}^{n+1/2}), \tag{4.2}$$

where $\mathcal{G}(\cdot)$ is specified in (2.3), that appears within Algorithm 3.1. To this end consider N = 3 and the velocities and Riemann initial data

$$\boldsymbol{v}^{\max} = (1,3,6)^{\mathrm{T}}, \quad \Phi(x,0) = \Phi_0(x) = \begin{cases} \Phi_{\mathrm{L}} = (0.05, 0.08, 0.12)^{\mathrm{T}} & \text{for } x < 0, \\ \Phi_{\mathrm{R}} = (0.14, 0.16, 0.12)^{\mathrm{T}} & \text{for } x \ge 0. \end{cases}$$



FIGURE 7. Example 5: numerical solution for a free-flow regime $(\mathcal{G}(t) = \alpha_V)$: (a) initial condition, (b, c) density profiles with M = 1600 at simulated time (b) T = 0.1, (c) T = 0.2.



FIGURE 8. Example 5: simulated total density computed with BCOV scheme with N = 3 and M = 1600: (a) free flow ($\mathcal{G}(t) = \alpha_V$), (b) congested flow ($\mathcal{G}(t) = 0$).

Observe that $\phi_{\rm R} = \phi^* = s(t)$, where $\phi_{\rm R}$ is the total density of the right state $\Phi_{\rm R}$.

As for Example 3 we show that the solution depends on the boundary condition $\mathcal{F}(t) \in (\mathbf{v}^{\max})^{\mathrm{T}} \mathbf{s}(t) \tilde{V}(\mathbf{s}(t))$. We start with the initial condition shown in Figure 7 (a) and compute the solution twice, once using $\mathcal{G}(t) = \alpha_V$, and the second time using $\mathcal{G}(t) = 0$. In Figures 7 (b) and (c) we display the profile for each class and total density for the first case $\mathcal{G}(t) = \alpha_V$ at two different simulated times. We can see that in this case a free-flow regime is produced, which is verified in Figure 8 (a). In Figure 9 we display the profiles for each class and total density for the second case $\mathcal{G}(t) = 0$ at two different simulation time. In contrast to the previous cases, a congested flow regime is produced, as is illustrated in Figure 8 (b).

4.6. Example 6: multiclass case (N = 5), smooth initial condition. In this example we consider N = 5, the velocities $v^{\max} = (1, 2, 3, 4, 5)^{\mathrm{T}}$, and the initial condition

 $\Phi(x,0) = \Phi_0(x) = (0.15, 0.2, 0.3, 0.2, 0.15)^{\mathrm{T}} \psi(x), \quad \psi(x) = \exp(-50(x+2)^2/3).$



FIGURE 9. Example 5: numerical solution for a congested flow regime ($\mathcal{G}(t) = 0$): density profiles with M = 1600 at simulated time (a) T = 0.1, (b) T = 0.2. The initial condition is the same as in Figure 7 (a).

	T = 0.02	T = 0.12
M	$e_M(\phi^{\Delta})$	$e_M(\phi^{\Delta})$
100	1.39e-2	3.87e-2
200	7.90e-3	2.47e-2
400	4.20e-3	1.55e-2
800	2.00e-3	9.20e-3
1600	1.00e-3	5.10e-3

TABLE 2. Example 6: approximate L^1 errors $e_M(u)$ with $\Delta x = 2/M$.

We display in Figure 10 numerical approximation computed with M = 1600 at simulation time T = 0.02 and T = 0.12. We observe the dynamics of each individual densities ϕ_i and the total density ϕ , which exhibits a shock wave due to the discontinuity in the flux. This behaviour is similar to that presented in Figure 4. In Figure 11 we display the evolution of $\phi^{\Delta}(\cdot, t)$ for $t \in [0, 0.12]$, and we compare the solution with the approximation of the continuous problem (where $\alpha_V = 0$). For the discontinuous case the shock is more clearly observed than in the continuous case. In Figures 12 and 13 we compare the numerical approximation computed with M = 100, with a reference solution at simulated times T = 0.02 and T = 0.12. In Table 1 we compute the approximate L^1 error based on a reference solution obtained by the BCOV scheme with $M_{\rm ref} = 12800$. We observe that the approximate L^1 errors decrease as the grid is refined.

4.7. Example 7: multiclass case (N = 5), bimodal smooth initial condition. In this example we consider N = 5, the velocity vector $\boldsymbol{v}^{\max} = (1, 1.5, 2, 6, 7)^{\mathrm{T}}$, and the initial condition

 $\Phi(x,0) = \Phi_0(x) = (0.17, 0.17, 0.16, 0, 0)^{\mathrm{T}} \psi_1(x) + (0, 0, 0, 0.245, 0.245)^{\mathrm{T}} \psi_2(x),$



FIGURE 10. Example 6: numerical solutions obtained with BCOV scheme with N = 5 and M = 1600 at simulated times (a) T = 0.02, (b) T = 0.12.



FIGURE 11. Example 6: simulated total density obtained with BCOV scheme with N = 5 and M = 1600: (a) discontinuous problem, (b) continuous problem.

where we define

 $\psi_1(x) = \exp(-10(x-2)^2), \quad \psi_2(x) = \exp(-50(x-1)^2/4)$

for $x \in [0, 5]$. We compute numerical approximation at simulated times T = 0.1, T = 0.2 and T = 0.3 with different discretizations by using $M = 100 \times 2^{l}$ and $l = 0, 1, \ldots, 4$. In Table 4.7 we compute the L^{1} error comparing with respect to a reference solution computed by the BCOV scheme with $M_{\rm ref} = 12800$. We observe that the approximate L^{1} errors decrease as the grid is refined. Again, this behaviour is similar to that observed in Figure 4. Figure 14 shows results for $M = M_{\rm ref} = 12800$. The numerical results of Figure 14 indicate that jumps in the total density ϕ only occur from smaller to higher values in an increasing x-direction.

MULTICLASS TRAFFIC WITH DISCONTINUOUS VELOCITY



FIGURE 12. Example 6: comparison of reference solution ($M_{\text{ref}} = 12800$) with approximate solutions computed by BCOV scheme with M = 100 at simulated time T = 0.02.

	T = 0.1	T = 0.2	T = 0.3
M	$e_M(\phi^{\Delta})$	$e_M(\phi^{\Delta})$	$e_M(\phi^{\Delta})$
100	7.42e-2	9.50e-2	1.06e-1
200	4.12e-2	5.50e-2	6.49e-2
400	2.27e-2	3.34e-2	3.88e-2
800	1.24e-2	1.97-2	2.35e-2
1600	6.50e-3	1.10e-2	1.35e-2

TABLE 3. Example 7: Approximate L^1 errors $e_M(u)$ with $\Delta x = 5/M$.

This phenomenon occurs because the speeds of the last two classes are greater than the first three. Furthermore, in Figure 15 we show the simulated total density computed by the BCOV scheme with N = 5 and M = 1600.

5. **Conclusions.** We have proposed a numerical scheme for a multiclass Lighthill-Whitham-Richards model with a velocity function that is discontinuous in the solution variable. The treatment is motivated and in part based on the numerical scheme proposed by Towers [28]. However, in contrast to that approach we have assume that the discontinuity is present in the velocity function (not in the flux); this observation makes it possible to and construct an alternative scheme based on Scheme 4 of [6]. Furthermore, we have seen that our scheme can be easily extended to the multiclass case. We have proved for the scalar case that the numerical approximations convergence to a weak solution and for the multiclass case that the



FIGURE 13. Example 6: comparison of reference solution ($M_{\rm ref} = 12800$) with approximate solutions computed by BCOV scheme with M = 100 at simulated time T = 0.12.



FIGURE 14. Example 7: numerical solution computed with BCOV scheme with N = 5 and M = 12800 at simulated times (a) T = 0.1, (b) T = 0.2 and (c) T = 0.3.

scheme preserves an invariant region. Examples 1 to 3 indicate that the scheme converges to the same weal solution as that of [28], and all numerical examples indicate that our scheme converges in both the scalar and multiclass cases.

The present analysis and numerical method can be extended in several directions. Concerning the model itself, at the moment a certain shortcoming is the limitation to the initial-boundary value problem on a fixed road segment. This is due to the particular boundary condition (1.5). It seems desirable to obtain a formulation for a closed road with periodic boundary conditions (a configuration that is commonly



FIGURE 15. Example 7: simulated total density computed with BCOV scheme with N = 5 and M = 1600.

studied in traffic modeling to analyze, say, the formation of stop-and-go waves; cf., e.g., [5, 11]). However, it is not obvious whether the way the boundary condition is posed allows "gluing together" the ends of the computational domain to create a "seamless" closed circuit. Open issues also include the incorporation of discontinuities in spatial position (akin to the treatment in [9]), and the discussion of the notion of entropicity. In fact, the issue of convergence to an entropy solution is an open problem even in the scalar case for both the scheme advanced in [28] as well as the present approach. Likewise, we recall that for general N the MCLWR model with a Lipschitz continuous function V admits a separable entropy function (see [1,2]) that can be utilized, for instance, to construct entropy stable schemes [10]. It remains to be explored whether these concepts are meaningful for the MCLWR model with a discontinuous velocity function V. Finally, it is clear that the numerical method is formally first-order accurate and can possibly improved by known techniques (e.g., weighted essentially non-oscillatory (WENO) reconstructions in combination with higher-order time integrators).

Acknowledgements. The authors are supported by the INRIA Associated Team "Efficient numerical schemes for non-local transport phenomena" (NOLOCO; 2018–2020). RB, RO and LMV also acknowledge support by project CMM, ANID/PIA/AFB170001. In addition, LMV is supported by Fondecyt project 1181511 and RB by projects Fondecyt 1170473 and CRHIAM, ANID/FONDAP/15130015.

References

- S. Benzoni-Gavage and R. M. Colombo, An n-populations model for traffic flow, Eur. J. Appl. Math. 14 (2003), 587–612.
- [2] S. Benzoni-Gavage, R. M. Colombo, and P. Gwiazda, Measure valued solutions to conservation laws motivated by traffic modelling, *Proc. Royal Soc. A*, 462 (2006), 1791–1803.
- [3] M. Bulíček, P. Gwiazda, J. Málek, and A. Świerczewska-Gwiazda, On scalar hyperbolic conservation laws with a discontinuous flux, *Math. Models Methods Appl. Sci.*, **21** (2011), 89–113.
- [4] M. Bulíček, P. Gwiazda, and A. Świerczewska-Gwiazda, Multi-dimensional scalar conservation laws with fluxes discontinuous in the unknown and the spatial variable, *Math. Models Methods Appl. Sci.*, 23 (2013), 407–439.

- [5] R. Bürger, C. Chalons, and L.M. Villada, Anti-diffusive and random-sampling Lagrangianremap schemes for the multi-class Lighthill-Whitham-Richards traffic model, *SIAM J. Sci. Comput.*, **35** (2013), B1341–B1368.
- [6] R. Bürger, A. García, K. H. Karlsen, and J. D. Towers, A family of numerical schemes for kinematic flows with discontinuous flux, J. Eng. Math, 60 (2008), 387–425.
- [7] R. Bürger, A. García, K.H. Karlsen, and J.D. Towers, Difference schemes, entropy solutions, and speedup impulse for an inhomogeneous kinematic traffic flow model, *Netw. Heterog. Media*, 3 (2008), 1–41.
- [8] R. Bürger, K. H. Karlsen, H. Torres, and J. D. Towers, Second-order schemes for conservation laws with discontinuous flux modelling clarifier-thickener units, *Numer. Math.*, **116** (2010), 579–617.
- [9] R. Bürger, K.H. Karlsen, and J.D. Towers, On some difference schemes and entropy conditions for a class of multi-species kinematic flow models with discontinuous flux, *Netw. Heterog. Media*, 5 (2010), 461–485.
- [10] R. Bürger, H. Torres, and C.A. Vega, An entropy stable scheme for the multiclass Lighthill-Whitham-Richards traffic model, Adv. Appl. Math. Mech., 11 (2019), 1022–1047.
- [11] R. Bürger, P. Mulet, and L. M. Villada, A diffusively corrected multiclass Lighthill-Whitham-Richards traffic model with anticipation lengths and reaction times, *Adv. Appl. Math. Mech.*, 5 (2013), 728–758.
- [12] J. Carrillo, Conservation law with discontinuous flux function and boundary condition, J. Evol. Equ., 3 (2003), 283–301.
- [13] C. Chalons and P. Goatin, Godunov scheme and sampling technique for computing phase transitions in traffic flow modeling, *Interf. Free Bound.*, 10 (2008), 197–221.
- [14] R.M. Colombo, Hyperbolic phase transitions in traffic flow, SIAM J. Appl. Math., 63 (2002), 708–721.
- [15] J. P. Dias and M. Figueira, On the Riemann problem for some discontinuous systems of conservation laws describing phase transitions, *Commun. Pure Appl. Anal.*, 3 (2004), 53–58.
- [16] J. P. Dias, M. Figueira, On the approximation of the solutions of the Riemann problem for a discontinuous conservation law, Bull. Braz. Math. Soc. New Ser., 36 (2005), 115–125.
- [17] J. P. Dias, M. Figueira, and J. F. Rodrigues, Solutions to a scalar discontinuous conservation law in a limit case of phase transitions, J. Math. Fluid Mech., 7 (2005), 153–163.
- [18] S. Diehl, A conservation law with point source and discontinuous flux function, SIAM J. Math. Anal., 56 (1996), 388–419.
- [19] R. Donat and P. Mulet, Characteristic-based schemes for multi-class Lighthill-Whitham-Richards traffic models, J. Sci. Comput., 37 (2008), 233–250.
- [20] R. Donat and P. Mulet, A secular equation for the Jacobian matrix of certain multi-species kinematic flow models, *Numer. Methods Partial Differential Equations*, 26 (2010), 159–175.
- [21] T. Gimse, Conservation laws with discontinuous flux functions, SIAM J. Numer. Anal., 24 (1993), 279–289.
- [22] T. Gimse and N. H. Risebro, Solution to the Cauchy problem for a conservation law with a discontinuous flux function, SIAM J. Math. Anal., 23 (1992), 635–648.
- [23] H. Holden and N. H. Risebro, Front Tracking for Hyperbolic Conservation Laws, Second Ed., Springer-Verlag, Berlin, 2015.
- [24] M.J. Lighthill and G.B. Whitham, On kinematic waves: II. A theory of traffic flow on long crowded roads, Proc. Royal Soc. A, 229 (1955), 317–345.
- [25] Y. Lu, S. Wong, M. Zhang, and C.-W. Shu, The entropy solutions for the Lighthill-Whitham-Richards traffic flow model with a discontinuous flow-density relationship, *Transp. Sci.*, 43 (2009), 511–530.
- [26] S. Martin and J. Vovelle, Convergence of the finite volume method for scalar conservation law with discontinuous flux function, ESAIM Math. Model. Numer. Anal., 42 (2008), 699–727.
- [27] P.I. Richards, Shock waves on the highway, Oper. Res., 4 (1956), 42–51.
- [28] J.D. Towers, A splitting algorithm for LWR traffic models with flux discontinuities in the unknown, J. Comput. Phys., 421 (2020), article 109722 (30pp).
- [29] J. K. Wiens, J. M. Stockie, and J. F. Williams, Riemann solver for a kinematic wave traffic model with discontinuous flux, J. Comput. Phys., 242 (2013), 1–23.
- [30] G. C. K. Wong and S. C. Wong, A multi-class traffic flow model—an extension of LWR model with heterogeneous drivers, *Transp. Res. A*, 36 (2002), 827–841.
- [31] P. Zhang, R. X. Liu, S. C. Wong, and D. Q. Dai, Hyperbolicity and kinematic waves of a class of multi-population partial differential equations, *Eur. J. Appl. Math*, **17** (2006), 171–200.

- [32] M. Zhang, C.-W. Shu, G. C. K. Wong, and S. C. Wong, A weighted essentially non-oscillatory numerical scheme for a multi-class Lighthill-Whitham-Richards traffic flow model, J. Comput. Phys., 191 (2003) 639–659.
- [33] P. Zhang, S. C. Wong, and C.-W. Shu, A weighted essentially non-oscillatory numerical scheme for a multi-class traffic flow model on an inhomogeneous highway, J. Comput. Phys., 212 (2006), 739–756.
- [34] P. Zhang, R.-X. Liu, S.C. Wong, and S.Q. Dai, Hyperbolicity and kinematic waves of a class of multi-population partial differential equations, *Eur. J. Appl. Math.*, **17** (2006), 171–200.
- [35] P. Zhang, S.C. Wong, and S.Q. Dai, A note on the weighted essentially non-oscillatory numerical scheme for a multi-class Lighthill-Whitham-Richards traffic flow model, *Commun. Numer. Meth. Eng.*, 25 (2009), 1120–1126.
- [36] P. Zhang, S.C. Wong, and Z. Xu, A hybrid scheme for solving a multi-class traffic flow model with complex wave breaking, *Comput. Methods Appl. Mech. Engrg.*, **197** (2008), 3816–3827.

E-mail address: rburger@ing-mat.udec.cl *E-mail address*: christophe.chalons@uvsq.fr *E-mail address*: rordonez@udec.cl *E-mail address*: lvillada@ubiobio.cl

Centro de Investigación en Ingeniería Matemática (CI²MA)

PRE-PUBLICACIONES 2020

- 2020-15 FELIPE LEPE, DAVID MORA, GONZALO RIVERA, IVÁN VELÁSQUEZ: A virtual element method for the Steklov eigenvalue problem allowing small edges
- 2020-16 JOSÉ QUERALES, RODOLFO RODRÍGUEZ, PABLO VENEGAS: Numerical approximation of the displacement formulation of the axisymmetric acoustic vibration problem
- 2020-17 PATRICK E. FARRELL, LUIS F. GATICA, BISHNU LAMICHHANE, RICARDO OYAR-ZÚA, RICARDO RUIZ-BAIER: Mixed Kirchhoff stress - displacement - pressure formulations for incompressible hyperelasticity
- 2020-18 DAVID MORA, IVÁN VELÁSQUEZ: Virtual elements for the transmission eigenvalue problem on polytopal meshes
- 2020-19 GABRIEL N. GATICA, GEORGE C. HSIAO, SALIM MEDDAHI: Further developments on boundary-field equation methods for nonlinear transmission problems
- 2020-20 NICOLAS BARNAFI, GABRIEL N. GATICA, DANIEL E. HURTADO, WILLIAN MI-RANDA, RICARDO RUIZ-BAIER: New primal and dual-mixed finite element methods for stable image registration with singular regularization
- 2020-21 GONZALO A. BENAVIDES, SERGIO CAUCAO, GABRIEL N. GATICA, ALEJANDRO A. HOPPER: A new non-augmented and momentum-conserving fully-mixed finite element method for a coupled flow-transport problem
- 2020-22 RAIMUND BÜRGER, ELVIS GAVILÁN, DANIEL INZUNZA, PEP MULET, LUIS M. VILLADA: Exploring a convection-diffusion-reaction model of the propagation of forest fires: computation of risk maps for heterogeneous environments
- 2020-23 GABRIEL N. GATICA, RICARDO OYARZÚA, RICARDO RUIZ-BAIER, YURI D. SO-BRAL: Banach spaces-based analysis of a fully-mixed finite element method for the steady-state model of fluidized beds
- 2020-24 JESSIKA CAMAÑO, SERGIO CAUCAO, RICARDO OYARZÚA, SEGUNDO VILLA-FU-ENTES: A posteriori error analysis of a momentum conservative Banach-spaces based mixed-FEM for the Navier-Stokes problem
- 2020-25 RAIMUND BÜRGER, JULIO CAREAGA, STEFAN DIEHL: A method-of-lines formulation for a model of reactive settling in tanks with varying cross-sectional area
- 2020-26 RAIMUND BÜRGER, CHRISTOPHE CHALONS, RAFAEL ORDOÑEZ, LUIS M. VILLADA: A multiclass Lighthill-Whitham-Richards traffic model with a discontinuous velocity function

Para obtener copias de las Pre-Publicaciones, escribir o llamar a: DIRECTOR, CENTRO DE INVESTIGACIÓN EN INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C, CONCEPCIÓN, CHILE, TEL.: 41-2661324, o bien, visitar la página web del centro: http://www.ci2ma.udec.cl









Centro de Investigación en Ingeniería Matemática (CI²MA) **Universidad de Concepción**

Casilla 160-C, Concepción, Chile Tel.: 56-41-2661324/2661554/2661316http://www.ci2ma.udec.cl





